



## <u>Do's and Don'ts for mOPV1 and mOPV3 Distribution</u> Regional Stores

The vaccine used during the coming polio campaign will be monovalent OPV type 1 or 3 (mOPV1 or mOPV3). MOPV is the most appropriate vaccine for the type of virus causing polio during the current outbreak. Trivalent OPV (tOPV) will not be used.

It is critical to prevent the use of tOPV during the campaign and the use of mOPV during the routine after the campaign.

mOPV should be stored and used under the same conditions as tOPV.

The role of the Regional Store is **<u>critical</u>** to ensuring that the right vaccine gets to the right place. Regional stores should follow these Do's and Don'ts:

piace.	Re	egional stores should follow these Do's and Don'ts:	✓
Do!	1.	Ensure that everyone working at the store who handles vaccine is aware of the 2 types of vaccine being used in the NID.	
	2.	In stock keeping records of polio vaccine, ensure mOPV is specified as opposed to tOPV.	
	3.	At Regional level, use separate deep-freezer or ice-lined refrigerator (ILR) for mOPV and tOPV.	
	4.	When mOPV vaccine is received, check each box to be sure it contains mOPV vials. Be careful – the 2 vaccine labels look very similar!	
		DUMMY  BELOW 1982  MICH TO ME PAULTINE  MICH TO ME PAULT NO  MICH TO ME PAULTINE  MICH TO ME PAULTINE  MICH TO ME PAULT NO  MICH	

- 5. Store mOPV in a separate deep-freezer / ILR, in the same conditions as usual. Clearly label as mOPV and post a list of districts to receive mOPV on the outside of the storage unit.
- 6. Label cold boxes for distribution to mOPV districts with a mOPV sticker.
- 7. Keep good records of the vaccine receipt and distribution.
- 8. If a district runs out of mOPV during the campaign, provide it with additional mOPV or arrange from a district with surplus mOPV supply.
- 9. Have all mOPV from the HC and (sub) districts return to the central store between the rounds, or ensure and check they keep their stocks rigorously separated.

## DON'T!

Don't use tOPV for the NID or mOPV for routine vaccination Don't store both vaccines in the same deep-freezer, ILR or cold box.

Don't distribute vaccine during the NID without checking carefully that it is mOPV.

Don't be afraid to ask - it's better to be sure than send the wrong vaccine to an area.