


MDT Regimens


Each blister pack contains treatment for 4 weeks.



PB adult treatment:

- Once a month:** Day 1
 – 2 capsules of rifampicin (300 mg X 2)
 – 1 tablet of dapsone (100 mg)
- Once a day:** Days 2–28
 – 1 tablet of dapsone (100 mg)
- Full course:** 6 blister packs

PB adult blister pack




MB adult treatment:

- Once a month:** Day 1
 – 2 capsules of rifampicin (300 mg X 2)
 – 3 capsules of clofazimine (100mg X 3)
 – 1 tablet of dapsone (100 mg)
- Once a day:** Days 2–28
 – 1 capsule of clofazimine (50 mg)
 – 1 tablet of dapsone (100 mg)
- Full course:** 12 blister packs

MB adult blister pack

It is crucial that patients understand which drugs they have to take once a month and which every day.




PB child treatment (10–14 years):

- Once a month:** Day 1
 – 2 capsules of rifampicin (300 mg+150 mg)
 – 1 tablet of dapsone (50 mg)
- Once a day:** Days 2–28
 – 1 tablet of dapsone (50 mg)
- Full course:** 6 blister packs

PB child blister pack

For children younger than 10, the dose must be adjusted according to body weight.



MB child treatment (10–14 years):

- Once a month:** Day 1
 – 2 capsules of rifampicin (300 mg+150 mg)
 – 3 capsules of clofazimine (50 mg X 3)
 – 1 tablet of dapsone (50 mg)
- Once a day:** Days 2–28
 – 1 capsule of clofazimine every other day (50 mg)
 – 1 tablet of dapsone (50 mg)
- Full course:** 12 blister packs

MB child blister pack

For children younger than 10, the dose must be adjusted according to body weight.