Philippines

Total population: 93 444 000 > Population aged 15 years and older (15+): 65% > Population in urban areas: 49% > Income group (World Bank): Lower middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2010 Data refer to litres of pure alcohol per capita (15+).



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	Average 2003—2005		rage -2010	Change
Recorded	4.4	4	.6	+
Unrecorded	2.0	0	.9	*
Total	6.4	5.4		*
Total males / females		9.2	1.7	
WHO Western Pacific Region	5.4	6.8		

Prevalence of heavy episodic drinking* (%), 2010

	Population	Drinkers only
Males (15+)	3.5	6.3
Females (15+)	0.3	0.8
Both sexes (15+)	1.9	4.2

*Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2012 $\,$

	ASDR*		AAF (%)	
Liver cirrhosis, males / females	35.0	9.7	66.7	49.6
Road traffic accidents, males / females	22.8	6.1	9.9	2.0
*Per 100 000 population (15+).				

Years of life lost (YLL) score*, 2012	LEAST < 1 2 3 4 5 > MOST
*Pacad on alaphal attributable years of life last	

*Based on alcohol-attributable years of life lost

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No /
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions for on-/off-premise sales of alcoholic beverages: Hours, days / places, density Specific events / intoxicated persons / petrol stations	Yes, No / Yes, No No / No / No

Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, $2010\,$



Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2010

Males (15+)	16.7
Females (15+)	5.2
Both sexes (15+)	12.3

Abstainers (%), 2010

	Males	Females	Both sexes
Lifetime abstainers (15+)	21.5	46.2	34.0
Former drinkers* (15+)	23.2	20.3	21.7
Abstainers (15+), past 12 months	44.7	66.5	55.7
*Persons who used to drink alcoholic beverages but have not done so in the past 12 months.			

Patterns of drinking score, 2010 LEAST RISKY < 1 2 3 4 5 > MOST RISKY

Prevalence of alcohol use disorders and alcohol dependence (%), 2010*

	Alcohol use disorders**	Alcohol dependence
Males	7.7	5.2
Females	1.4	0.7
Both sexes	4.6	2.9
WHO Western Pacific Region	4.6	2.3

*12-month prevalence estimates (15+).

**Including alcohol dependence and harmful use of alcohol.

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in $\%$	0.05 / 0.05 / 0.05
Legally binding regulations on alcohol advertising / product placement	No / No
Legally binding regulations on alcohol sponsorship / sales promotion	No / No
Legally required health warning labels on alcohol advertisements / containers	Yes / Yes
National government support for community action	No
National monitoring system(s)	No

© World Health Organization 2014