

[Back to UNICEF.org](#)



UNICEF Data: Monitoring the Situation of Children and Women

ABOUT US

RESOURCES

CONTACT

Home / [NAM](#)

STATISTICS BY TOPIC + COUNTRY

MULTIPLE INDICATOR CLUSTER SURVEYS (MICS)

NAMIBIA

NAMIBIA

KEY DEMOGRAPHIC INDICATORS

Under 5 Mortality Indicator bubble

45

PER 1,000 LIVE BIRTHS

Under 5 Mortality Indicator bubble

2,459,000



TRENDS IN UNDER-FIVE MORTALITY RATE

Under-five mortality rate

Deaths per 1,000 live births

Visit childmortality.org for complete data

DISPARITIES BY HOUSEHOLD WEALTH

Oral Rehydration:

Children under five with diarrhoea receiving ORS (%)

Secondary education:

Net attendance ratio in secondary education (%)

Birth Registration:

Children under age 5 whose births are registered (%)

Visit devinfo.org/unicefdata for complete database

CHILD HEALTH

Proportion of under-five children with suspected pneumonia taken to health provider (%)

68

Proportion of children under five years old with diarrhoea receiving oral rehydration salts (%)

72

Proportion of children under five sleeping under insecticide-treated nets (%)

6

Proportion of households owning at least one insecticide-treated net (ITN) (%)

24

CHILD MORTALITY

Under-five mortality rate (U5MR), deaths per 1,000 live births

45

Number of under-five deaths

3,318

Infant mortality rate (IMR), deaths per 1,000 live births

33

Neonatal mortality rate (NMR), deaths per 1,000 live births

16

Proportion of 1 year-old children fully immunised against DPT (%)

92

Proportion of 1 year-old children immunised against measles (%)

85

[Read More About Child Health](#)

Under-five mortality rate (U5MR), deaths per 1,000 live births (male)

49

Under-five mortality rate (U5MR), deaths per 1,000 live births (female)

41

[Read More About Child Mortality](#)

NUTRITION

Early initiation of breastfeeding (within one hour of birth) (%)

71

Exclusive breastfeeding rate (%)

49

Continued breastfeeding rate at one year (%)

64

Prevalence of moderate and severe stunting (%)

23

Vitamin A supplementation (full) (%)

62

Proportion of households consuming iodized salt (%)

MATERNAL HEALTH

Proportion of women aged 15-49 who received postnatal care within 2 days after giving birth (%)

69

Antenatal care coverage for at least one visit (%)

97

Antenatal care coverage for at least four visits (%)

63

Proportion of births attended by skilled health personnel (%)

89

Caesarean section (%)

14

Proportion of women 20-24 years old who gave birth before age 18 (%)

57[Read More About Child Nutrition](#)**15**[Read More About Maternal Health](#)**EARLY CHILDHOOD**

Attendance in early childhood education(%)

..

Support for learning (any household member) (%)

..

Support for learning (father) (%)

..

Learning materials at home – children's books (%)

..

Learning materials at home – playthings (%)

..

Children left in inadequate care (%)

..

[Read More About Early Childhood Development](#)**EDUCATION**

Gross enrollment ratio (GER) in pre-primary education (%)

16

Net attendance ratio in primary education (NAR) (%)

92

Proportion of pupils starting grade 1 who reach grade 5 (%)

85

Proportion of out-of-school children of primary school age (%)

8

Net attendance ratio in secondary education (%)

60

Literacy rate (15-24 years) (%)

87[Read More About Education](#)**HIV/AIDS****CHILD PROTECTION**

Percentage of pregnant women living with HIV receiving most effective ARVs for PMTCT (%)

95

Estimated number of new HIV infections among children 0-14 years

<500

HIV positive children (aged 0-14) receiving antiretroviral therapy (%)

>95

Adolescent girls aged 15-19 who were tested for HIV in the last 12 months and received the results (%)

29

Adolescent boys aged 15-19 who were tested for HIV in the last 12 months and received the results (%)

14

Estimated number of new HIV infections among adolescents aged 15-19

<1000

[Read More About HIV/AIDS](#)

Children under age 5 whose birth is registered (%)

87

Women aged 20-24 years who were first married or in union by age 18 (%)

7

Children aged 5-14 years engaged in child labour (%)

..

Children aged 2 to 14 years who experienced any violent discipline (psychological aggression and/or physical punishment) in the past month (%)

..

Justification of wife beating (female) (%)

28

Justification of wife beating (male) (%)

22

[Read More About Child Protection](#)

SANITATION

Proportion of population using improved sanitation facilities (%)

34

Proportion of population using shared sanitation facilities (%)

WATER

Proportion of population using a piped drinking water supply on premises (%)

51

Proportion of population using another improved drinking water source (%)

12

Proportion of population using other unimproved sanitation facilities (%)

6

Proportion of population practicing open defecation (%)

48[Read More About Sanitation](#)**40**

Proportion of population using an unimproved drinking water source other than surface water (%)

1

Proportion of population using a surface water source (%)

8[Read More About Water](#)**ADDITIONAL RESOURCES**[Namibia](#)**MICS**[Namibia](#)[About UNICEF](#) [Contact Us](#) [Legal](#)