Total population: 28 514 000

Income group: Low

fomales

## Mortality\*

#### Number of diabetes deaths

|            | Illaics | Terriares |
|------------|---------|-----------|
| ages 30–69 | 1 270   | 1 080     |
| ages 70+   | 1 370   | 1 430     |

malac


fomales


#### Number of deaths attributable to high blood glucose

|            | marcs | Terriares |
|------------|-------|-----------|
| ages 30–69 | 2 980 | 1 970     |
| ages 70+   | 2 720 | 2 590     |

# Proportional mortality (% of total deaths, all ages)\*

# Trends in age-standardized prevalence of diabetes





## Prevalence of diabetes and related risk factors

|                     | males | females | total |
|---------------------|-------|---------|-------|
| Diabetes            | 10.5% | 7.9%    | 9.1%  |
| Overweight          | 13.6% | 19.6%   | 16.7% |
| Obesity             | 1.7%  | 4.1%    | 2.9%  |
| Physical inactivity | 3.9%  | 2.8%    | 3.3%  |

## National response to diabetes

### Policies, guidelines and monitoring

| Operational policy/strategy/action plan for diabetes                                 | Yes           |
|--|---------------|
| Operational policy/strategy/action plan to reduce overweight and obesity             | No            |
| Operational policy/strategy/action plan to reduce physical inactivity                | Yes           |
| Evidence-based national diabetes guidelines/protocols/standards                      | Not available |
| Standard criteria for referral of patients from primary care to higher level of care | NR            |
| Diabetes registry  | No            |
| Recent national risk factor survey in which blood glucose was measured               | Yes           |

## Availability of medicines, basic technologies and procedures in the public health sector

#### Medicines in primary care facilities

| Insulin                                      | O |
|--|---|
| Metformin                                    | • |
| Sulphonylurea                                | 0 |
| Procedures                                   |   |
| Retinal photocoagulation                     | 0 |
| Renal replacement therapy by dialysis        | 0 |
| Renal replacement therapy by transplantation | • |

| blood glucose measurement                | O |
|--|---|
| Oral glucose tolerance test              | 0 |
| HbA1c test                               | 0 |
| Dilated fundus examination               | 0 |
| Foot vibration perception by tuning fork | 0 |

0

Basic technologies in primary care facilities

Urine strips for glucose and ketone measurement

Foot vascular status by Doppler

NR = country responded to survey but did not give a response to a specific question

World Health Organization - Diabetes country profiles, 2016.

<sup>\*</sup> The mortality estimates for this country have a high degree of uncertainty because they are not based on any national NCD mortality data (see Explanatory Notes).

<sup>○ =</sup> not generally available ■ = generally available