Liberia HIV Country Profile: 2017

WHO/CDS/HIV/18.50

Demographic and socioeconomic data

*** \$	4.7 million Total population (2017) UNPOP 710 US\$ GNI per capita, PPP (2017) World Bank
Q	725 Maternal mortality per 100 000 live births (2015) World Health Stat., WHO
\$	9.62% Health expenditure, total (% World Bank
∞\$	63 years Life expectancy at birth (20 UNPOP 68 US\$ Health expenditure per capi World Bank
	4.8 Total fertility rate (births pe UNPOP 0.44 Human Development Index UNPOP






Estimated ART coverage children and adults



Estimated % of pregnant women living with HIV who received ARVs for PMTCT


90-90-90 progress towards 2020 targets




Health sector cascade

	Value	Percent of 90-90-90
People living with HIV who know their status	N/A	N/A
Reported number of people living with HIV receiving ART	11 600	29%
People living with HIV with viral load suppression	N/A	N/A





Estimated number of deaths due to AIDS




Prevention of mother-to-child transmission (PMTCT)	Value	Year
Reported number of pregnant women living with HIV who received antiretrovirals for PMTCT	1 600	2017
Estimated percentage of pregnant women living with HIV who received antiretrovirals for PMTCT* ²	86%	2017
Final transmission rate including breastfeeding period*3	13.2	2017
Antenatal care coverage - at least one visit (%)	N/A	
Antenatal care coverage - at least four visits (%)	78%	2013
% of pregnant women with known HIV status	N/A	
Antenatal care attendees who were tested for syphilis (%)	8%	2015
Antenatal care attendees who tested positive for syphilis (%)	7.1%	2015
Antenatal care attendees who tested positive for syphilis and who received benzathine penicillin IM (%)	100%	2015
% of infants born to women living with HIV receiving a virological test within two months of birth (EID)* ¹	N/A	

^{*} Tracing indicators for elimination of monther-to-child transmission. A color scheme (green, orange,red) is used to indicate if the country is on target to meet 2020 targets: ¹90% ²95% ³Less than 5

Key populations

Percentage condom use among sex workers with most recent client	N/A
Needles distributed per person who inject drugs per year	N/A
HIV prevalence among men who have sex with men (%)	20%
HIV prevalence among sex workers (%)	N/A
HIV prevalence among people who inject drugs (%)	
Percentage of people who inject drugs receiving OST	N/A

Median annual price of ARV drug regimens for adults



National HIV policies and plans

Implementation of national policy on HIV self- testing (HIVST)	Policy on HIVST in development
Treatment initiation threshold adults and adolescents	≤500 cells/mm3 ●
Treatment initiation threshold children	Treat all, regardless of age
Implementation of treat all adults and adolescents	Recommending Treat all later in 2018

Dolutegravir (DTG) as part of 1st line ARVs for adults and adolescents	TDF/[3TC or FTC]/EFV first line ARVs only
Implementation of lifelong ART to pregnant & breastfeeding women	Implemented in a large number (>50%) of MCH sites
Implementation of national policy on viral load monitoring	Partially implemented
Completion of HIV services cascades	Data not available
Adaptation of WHO key population guidelines in national plans	No
National M&E plan or strategy for HIV	Yes, and updated in the past two years
Method to de-duplicate key data (i.e. unique identifiers)	Yes
Nationally representative quality assessment of facility-level ART data	Yes

 $\textcircled{\sc c}$ WHO 2018. Some rights reserved. Licence CC BY-NC-SA 3.0 IGO