

HIV and AIDS Estimates

Adults and children living with HIV	9200 [6800 - 12 000]
Adults aged 15 and over living with HIV	9100 [6800 - 12 000]
Women aged 15 and over living with HIV	2800 [2100 - 3700]
Men aged 15 and over living with HIV	6300 [4700 - 8200]
Children aged 0 to 14 living with HIV	<200 [<100 - <200]
Adult aged 15 to 49 HIV prevalence rate	0.1 [0.1 - 0.2]
Women aged 15 to 49 HIV prevalence rate	<0.1 [<0.1 - 0.1]
Men aged 15 to 49 HIV prevalence rate	0.2 [0.1 - 0.3]
HIV prevalence among young women	<0.1 [<0.1 - <0.1]
HIV prevalence among young men	<0.1 [<0.1 - <0.1]
Adults and children newly infected with HIV	<1000 [<1000 - 1300]
Adults aged 15 and over newly infected with HIV	<1000 [<1000 - 1300]
Women aged 15 and over newly infected with HIV	<500 [<200 - <500]
Men aged 15 and over newly infected with HIV	<1000 [<500 - <1000]
Children aged 0 to 14 newly infected with HIV	<100 [<100 - <100]
HIV incidence per 1000 population (adults 15-49)	0.16 [0.11 - 0.22]
HIV incidence per 1000 population (all ages)	0.10 [0.06 - 0.14]
Adult and child deaths due to AIDS	<500 [<200 - <500]
Deaths due to AIDS among adults aged 15 and over	<500 [<200 - <500]
Deaths due to AIDS among women aged 15 and over	<200 [<100 - <200]



Orphans due to AIDS aged 0 to 17	1800 [1000 - 2700]
Deaths due to AIDS among children aged 0 to 14	<100 [<100 - <100]
Deaths due to AIDS among men aged 15 and over	<200 [<200 - <500]

HIV testing and treatment cascade

People living with HIV	9200 [6800 - 12 000]	Source: Spectrum
People living with HIV who know their status	5400	Source: GAM (case surveillance)
Percent of people living with HIV who know their status	58 [43 - 76]	
People living with HIV who are on ART	2800	Source: Spectrum
Percent of people living with HIV who are on ART	30	
People living with HIV who have suppressed viral loads	1700	Source: GAM; Viral load testing coverage: 72%
Percent of people living with HIV who have suppressed viral loads	19	

Antiretroviral therapy (ART)

Coverage of adults and children receiving ART (%)	30 [22 - 40]
Adults aged 15 and over receiving ART	30 [22 - 39]
Women aged 15 and over receiving ART	22 [16 - 29]
Men aged 15 and over receiving ART	33 [25 - 44]
Children aged 0 to 14 receiving ART	76 [50 - >95]



Country factsheets

AZERBAIJAN | 2016

Number of adults and children receiving ART (#)	2800
Adults aged 15 and over receiving ART	2700
Women aged 15 and over receiving ART	620
Men aged 15 and over receiving ART	2100
Children aged 0 to 14 receiving ART	80
Late HIV diagnosis [with the initial CD4 cell count <200 cells/ mm3] (%)	21
Late HIV diagnosis among adults aged 15 and over	22
Late HIV diagnosis among women	15
Late HIV diagnosis among men	25
Adults and children newly-initiating ART (#)	578
Adults aged 15 and over newly-initiating ART	562
Children aged 0-14 newly-initiating ART	16
Women newly-initiating ART	202
Men newly-initiating ART	376
Adults and children known to be on ART 12 months after starting (%)	94
Adults known to be on ART 12 months after starting	94
Children known to be on ART 12 months after starting	>95

Elimination of mother-to-child transmission

Coverage of pregnant women who receive ARV for PMTCT (%)	75 [53 - >95]
Pregnant women who received ARV for PMTCT (#)	70
Pregnant women needing ARV for PMTCT (#)	<100 [<100 - <200]

Country factsheets

AZERBAIJAN | 2016

Early infant diagnosis (%)	61 [44 - 86]
New HIV infections averted due to PMTCT (%)	<100 [<100 - <100]

Sex workers

Population size estimate (#)	25 054	Source: GAM 2016; 2011 est.
HIV prevalence (%)	2.3	Source: Sentinel surveillance, 2016
Knowledge of HIV status (%)	37.2	Source: Behavioural Surveillance Survey, 2016
Condom use (%)	53	Source: « A A; 5 4 > 2 0 = 8 5 @ 0 A ? @ > A B @ 0 = 5 = > A B 8 '-8 = D 5 : F 8 8 8 D 0 : B > @ > 2 @ 8 A : 0 7 0 @ 0 6 5 = 8 O 2 ? > 2 5 4 5 = 8 8 = 5 : > B > @ K E C O 7 2 8 < K E 3 @ C ? ? = 0 A 5; 5 = 8 O 2 7 5 @ 1 0 9 4 6 0 = A : > 9 5 A ? C 1; 8 : 5 », 2011
Active syphilis (%)	11.2	Source: Sentinel surveillance 2016
Hepatitis B and coinfection with HIV (%)	8.7	Source: Behavioural Surveillance Survey, 2016
Hepatitis C and coinfection with HIV (%)	6	Source: Behavioural Surveillance Survey, 2016

Men who have sex with men



Population size estimate (#)	6572	Year of estimation: 2011
HIV prevalence (%)	2.2	Source: Sentinel surveillance 2016
Knowledge of HIV status (%)	69.7	Source: Behavioural Surveillance Survey, 2016
Condom use (%)	63.9	Source: Behavioural Surveillance Survey, 2016
Active syphilis (%)	13.6	Source: Sentinel surveillance 2016
Hepatitis B and coinfection with HIV (%)	7.2	Source: Behavioural Surveillance Survey, 2016
Hepatitis C and coinfection with HIV (%)	4	Source: Behavioural Surveillance Survey, 2016

People who inject drugs

Population size estimate (#)	71 283	Year of estimation: 2011
HIV prevalence (%)	8.5	Source: Sentinel surveillance 2016
Knowledge of HIV status (%)	12.2	Source: Behavioural Surveillance Survey, 2016
Condom use (%)	15.1	Source: Behavioural Surveillance Survey, 2016



Safe injecting practices (%)	46.3	Source: « A A; 5 4 > 2 0 = 8 5 @ 0 A ? @ > A B @ 0 = 5 = > A B 8 '-8 = D 5: F 8 8 8 D 0: B > @ > 2 @ 8 A: 0 7 0 @ 0 6 5 = 8 O 2 ? > 2 5 4 5 = 8 8 = 5: > B > @ K E C O 7 2 8 < K E 3 @ C?? = 0 A 5; 5 = 8 O 2 7 5 @ 1 0 9 4 6 0 = A: > 9 5 A ? C 1; 8: 5 », 2011
Needles and syringes distributed per person who injects drugs (#)	34.371	
Coverage of opioid substitution therapy	0.5	Source: National Programme data 2016
Hepatitis B and coinfection with HIV (%)	7.3	Source: Behavioural Surveillance Survey, 2016
Hepatitis C and coinfection with HIV (%)	43.9	Source: Behavioural Surveillance Survey, 2016

Prisoners

HIV prevalence (%)	2.8	Source: IBBS 2016
HIV prevention programmes in prisons: condoms distributed (#)	48 942	Source: Routine programme data 2016
Prevalence of TB infection among prisoners (%)	0.2	Representative of HIV/TB; Source: Routine programme data 2016



Prevalence of hepatitis C infection among prisoners (%)	2.2	Representative of HIV/HCV; Source: Routine programme data 2016
Hepatitis B and coinfection with HIV (%)	5.8	Source: Behavioural Surveillance Survey, 2016
Hepatitis C and coinfection with HIV (%)	25.8	Source: Behavioural Surveillance Survey, 2016

Combination prevention

Percentage of men 15-49 that are	96.9	Source: DHS 2006
circumcised (%)		

Gender

Prevalence of recent intimate partner violence among women aged 15-49 (%)	9.8	Source: 2006 DHS
Prevalence of recent intimate partner violence among women aged 15-19 (%)	11.5	Source: 2006 DHS
Prevalence of recent intimate partner violence among women aged 20-24 (%)	9.9	Source: 2006 DHS
Prevalence of recent intimate partner violence among women aged 25-49 (%)	9.8	Source: 2006 DHS

Young people



Country factsheets

AZERBAIJAN | 2016

Knowledge about HIV prevention among young people aged 15-24	14.9	Source: DHS 2011
Knowledge about HIV prevention among young women aged 15-24	14.9	Source: DHS 2011
Knowledge about HIV prevention among young men aged 15-24		

TB and HIV

Co-management of TB and HIV treatment (%)	61 [42 - 94]
Proportion of people living with HIV newly enrolled in HIV care with active TB disease (%)	8
Proportion of people living with HIV newly enrolled in HIV care started on TB preventive therapy (%)	60
Estimated TB-related deaths among people living with HIV (#)	<100 [<100 - <100]