INAIDS AIDSinfo

Country factsheets BOSNIA AND HERZEGOVINA | 2016

HIV and AIDS Estimates

Adults and children living with HIV	[]
Adults aged 15 and over living with HIV	[]
Women aged 15 and over living with HIV	[]
Men aged 15 and over living with HIV	[]
Children aged 0 to 14 living with HIV	[]
Adult aged 15 to 49 HIV prevalence rate	[]
Women aged 15 to 49 HIV prevalence rate	[]
Men aged 15 to 49 HIV prevalence rate	[]
HIV prevalence among young women	[]
HIV prevalence among young men	[]
Adults and children newly infected with HIV	[]
Adults aged 15 and over newly infected with HIV	[]
Women aged 15 and over newly infected with HIV	[]
Men aged 15 and over newly infected with HIV	[]
Children aged 0 to 14 newly infected with HIV	[]
HIV incidence per 1000 population (adults 15-49)	[]
HIV incidence per 1000 population (all ages)	[]
Adult and child deaths due to AIDS	[]
Deaths due to AIDS among adults aged 15 and over	[]
Deaths due to AIDS among women aged 15 and over	[]

UNAIDS | AIDSinfo

Country factsheets BOSNIA AND HERZEGOVINA | 2016

Deaths due to AIDS among men aged 15 and over	[]
Deaths due to AIDS among children aged 0 to 14	[]
Orphans due to AIDS aged 0 to 17	[]

HIV testing and treatment cascade

People living with HIV	<200 [<200 - <500]	Source: GAM (country supplied estimate)
People living with HIV who are on ART	<200	Source: GAM
Percent of people living with HIV who are on ART	81	

Antiretroviral therapy (ART)

Coverage of adults and children receiving ART (%)	[]
Adults aged 15 and over receiving ART	[]
Women aged 15 and over receiving ART	[]
Men aged 15 and over receiving ART	[]
Children aged 0 to 14 receiving ART	[]
Number of adults and children receiving ART (#)	•••
Adults aged 15 and over receiving ART	
Women aged 15 and over receiving ART	
Men aged 15 and over receiving ART	
Children aged 0 to 14 receiving ART	
Late HIV diagnosis [with the initial CD4 cell count <200 cells/ mm3] (%)	37

UNAIDS | AIDSinfo

Country factsheets BOSNIA AND HERZEGOVINA | 2016

Late HIV diagnosis among adults aged 15 and over	
Late HIV diagnosis among women	
Late HIV diagnosis among men	
Adults and children known to be on ART 12 months after starting (%)	83
Adults known to be on ART 12 months after starting	82
Children known to be on ART 12 months after starting	>95

Elimination of mother-to-child transmission

Coverage of pregnant women who receive ARV for PMTCT (%)	[]
Pregnant women who received ARV for PMTCT (#)	
Pregnant women needing ARV for PMTCT (#)	[]
New HIV infections averted due to PMTCT (%)	[]

Sex workers

Population size estimate (#)	4000	Source: GAM 2016; 2012 est.
HIV prevalence (%)		Source: VCCT reports 2016
Condom use (%)	80.4	Source: Behavioural Surveillance Survey, 2015

Men who have sex with men

UNAIDS | AIDSinfo

Country factsheets BOSNIA AND HERZEGOVINA | 2016

Population size estimate (#)	690	Year of estimation: 2012
HIV prevalence (%)	1.1	Source: VCCT 2016
Condom use (%)	62.4	Source: Behavioural Surveillance Survey, 2015

People who inject drugs

Population size estimate (#)	12 500	Year of estimation: 2012
HIV prevalence (%)		Source: VCCT reports 2016
Condom use (%)	35.2	Source: Behavioural Surveillance Survey, 2015
Safe injecting practices (%)	92.7	Source: Behavioural Surveillance Survey, 2015
Needles and syringes distributed per person who injects drugs (#)	142.128	
Coverage of opioid substitution therapy	11.3	Source: National Programme data 2016

Prisoners

HIV prevention programmes in prisons: condoms distributed (#)	20 232	Source: Routine programme data 2016
HIV prevention programmes in prisons: prisoners receiving opioid substitution therapy (#)	50	Source: Routine programme data 2016

Stigma and discrimination

Country factsheets BOSNIA AND HERZEGOVINA | 2016

Percentage of adults (15-49) who responded No to the question: Would you buy fresh vegetables from a shopkeeper or vendor if you knew that this person had HIV?	57.1	Source: 2011-12 MICS
Percentage of adults (15-49) who responded No to the question: Do you think that children living with HIV should be able to attend school with children who are HIV negative?		
Percentage of adults (15-49) who responded No to both questions		

TB and HIV

Co-management of TB and HIV treatment (%)	44 [31 - 68]
Proportion of people living with HIV newly enrolled in HIV care with active TB disease (%)	>95
Proportion of people living with HIV newly enrolled in HIV care started on TB preventive therapy (%)	>95
Estimated TB-related deaths among people living with HIV (#)	<100 [<100 - <100]