



Violence against women during COVID-19

15 April 2020 | Q&A

Home is not a safe place for me. What can I do?

I need medical attention because of violence. Who can help me and where should I go?

I am safe, but I am suffering from mental / sexual / social / long-term physical health problems because of violence. Is there anyone who can help me during COVID-19?

I am worried about someone I know. How can I help?

I want to report an incidence of violence during COVID-19. What should I do?

I have harmed or am worried about harming or hurting my partner (and children) with my words or actions. How can I stop?

I am a health worker. How can I help women survivors of violence during COVID-19?

I run a health facility. What can I do to support women survivors of violence during COVID-19?

I am a policy maker. What can I do to prevent and address violence against women during COVID-19?

Has violence against women increased since the beginning of the COVID-19 pandemic?

How does COVID-19 increase risks of violence for women?

Who is most vulnerable?

Why should I care about violence against women during COVID-19?

WHO TEAM Sexual and Reproductive Health and Research

Related

[Q&A on coronaviruses \(COVID-19\)](#)

[Q&A on COVID-19, pregnancy, childbirth and breastfeeding](#)

[Q&A on COVID-19, HIV and antiretrovirals](#)

[Q&A: Similarities and differences – COVID-19 and influenza](#)

[Q&A on Mass Gatherings and COVID-19](#)

[Q&A on infection prevention and control for health care workers caring for patients with suspected or confirmed 2019-nCoV](#)

[Q&A on COVID-19 and food and agriculture](#)

[Q&A on smoking and COVID-19](#)

[Q&A: Be active during COVID-19](#)

[Q&A on Malaria and the COVID-19 pandemic](#)

[Q&A: Contraception/family planning and COVID-19](#)

Fact sheets

[Violence against women](#)

29 November 2017

[Subscribe to the WHO newsletter →](#)