


Delivering mental health and psychosocial support interventions to women and children in conflict settings: a systematic review

Mahdis Kamali,¹ Mariella Munyuzangabo,¹ Fahad J Siddiqui,^{1,2} Michelle F Gaffey,¹ Sarah Meteke,¹ Daina Als,¹ Reena P Jain ,¹ Amruta Radhakrishnan,¹ Shailja Shah,¹ Anushka Atallahjan,¹ Zulfiqar A Bhutta^{1,3}

To cite: Kamali M, Munyuzangabo M, Siddiqui FJ, *et al*. Delivering mental health and psychosocial support interventions to women and children in conflict settings: a systematic review. *BMJ Global Health* 2020;**5**:e002014. doi:10.1136/bmjgh-2019-002014

Handling editor Seye Abimbola

▶ Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjgh-2019-002014>).

Received 20 September 2019
Revised 9 December 2019
Accepted 22 December 2019



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

¹Centre for Global Child Health, Hospital for Sick Children, Toronto, Ontario, Canada

²Health Services and Systems Research, Duke-NUS Graduate Medical School, Singapore, Singapore

³Center of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan

Correspondence to

Dr Zulfiqar A Bhutta;
Zulfiqar.bhutta@sickkids.ca

ABSTRACT

Background Over 240 million children live in countries affected by conflict or fragility, and such settings are known to be linked to increased psychological distress and risk of mental disorders. While guidelines are in place, high-quality evidence to inform mental health and psychosocial support (MHPSS) interventions in conflict settings is lacking. This systematic review aimed to synthesise existing information on the delivery, coverage and effectiveness of MHPSS for conflict-affected women and children in low-income and middle-income countries (LMICs).

Methods We searched Medline, Embase, Cumulative Index of Nursing and Allied Health Literature (CINAHL) and Psychological Information Database (PsycINFO) databases for indexed literature published from January 1990 to March 2018. Grey literature was searched on the websites of 10 major humanitarian organisations. Eligible publications reported on an MHPSS intervention delivered to conflict-affected women or children in LMICs. We extracted and synthesised information on intervention delivery characteristics, including delivery site and personnel involved, as well as delivery barriers and facilitators, and we tabulated reported intervention coverage and effectiveness data.

Results The search yielded 37 854 unique records, of which 157 were included in the review. Most publications were situated in Sub-Saharan Africa (n=65) and Middle East and North Africa (n=36), and many reported on observational research studies (n=57) or were non-research reports (n=53). Almost half described MHPSS interventions targeted at children and adolescents (n=68). Psychosocial support was the most frequently reported intervention delivered, followed by training interventions and screening for referral or treatment. Only 19 publications reported on MHPSS intervention coverage or effectiveness.

Discussion Despite the growing literature, more efforts are needed to further establish and better document MHPSS intervention research and practice in conflict settings. Multisectoral collaboration and better use of existing social support networks are encouraged to increase reach and sustainability of MHPSS interventions.

PROSPERO registration number CRD42019125221.

Key questions

What is already known about this topic?

- ▶ Armed conflicts are linked to mental health consequences that disproportionately burden vulnerable populations such as women and children, though empirical evidence on the mechanisms of intervention implementation and delivery is lacking.

What are the new findings?

- ▶ The majority of interventions delivered to children and adolescents were delivered in specialised centres and schools.
- ▶ School-based interventions appear to be effective at improving mental health outcomes in children and adolescents.
- ▶ Interventions targeted to women exclusively are very limited.
- ▶ Access and security to target populations, language and culture and inadequate infrastructure are some reported barriers to delivering mental health interventions.

Recommendations for policy

- ▶ Along with strengthening the documentation of research and practice of intervention delivery in conflict settings, more concerted efforts are needed to incorporate these interventions into established networks and resources.

BACKGROUND

Globally, 172 million people are affected by armed conflict, and at least 240 million children live in countries affected by conflict or fragility.^{1 2} Humanitarian crises, such as armed conflicts, are linked to increased levels of psychological distress and higher risk of an array of mental disorders.^{3–5} Complexities further increase when unstable situations result in large-scale displacement and migration, exposing populations to stressful events prior to departure, during transit and after

arrival, as they cope with integrating in a new environment or country.⁶ The mental health consequences of armed conflict are disproportionately experienced by children, who are already undergoing rapid and complex physiological, cognitive and emotional changes.⁷ Exposure to war, conflict and terror interferes with children's natural development and has shown to lead to elevated levels of post-traumatic stress, anxiety, depression and many behavioural and emotional reactions.^{7,8}

Empirical evidence on the delivery of effective mental health and psychosocial support (MHPSS) interventions to the most vulnerable populations, women and children, in humanitarian settings is scant. The 2017 Lancet series on health in humanitarian crises confirmed the dire necessity for vigorous, high quality and practical evidence to help inform interventions in these settings.⁹ The scarcity of empirical evidence is coupled with a lack of consensus on terminology, measurement and monitoring and evaluation of MHPSS interventions.¹⁰ As an attempt to safeguard best practices for MHPSS in emergency settings, WHO initiated the Inter-Agency Committee Guidelines for MHPSS in emergency settings in 2007.¹¹ They provide standards on implementing minimum, yet essential, responses as quickly as possible during emergencies. Additionally, the Sphere Handbook has become a reference tool on universal minimum standards for humanitarian response, but a particular focus on MHPSS is lacking.¹²

Previous literature reviews and evidence syntheses have been conducted on the prevalence of mental health disorders among conflict-affected persons,^{5,8,13–22} but these were limited to certain mental health conditions, focused on high-income settings, or are not up-to-date. Stronger scientific evidence on implementing effective MHPSS interventions is needed in order to adequately and effectively respond to humanitarian crises. This current systematic review aims to help fill this gap by synthesising existing data and information on how MHPSS interventions are being implemented and delivered in conflict settings, and examining the reported coverage and observed effects of these interventions on mental health outcomes.

METHODS

Indexed literature search

We conducted a systematic search of literature published between January 1 1990 and 31 March 2018 in Medline, Embase, CINAHL and PsycINFO using search terms related to three concepts: (1) conflict; (2) women and children; and (3) MHPSS. Conflict-related terms included “war”, “crisis”, “refugees”, “internally displaced person (IDP)” and “stateless”. Population-related words included “women”, “children”, “pregnant”, “adolescents”, and “newborn”. MHPSS-related terms included “mental disorders”, “mental illness”, “mental symptoms”, “psychiatric diagnoses”, and so on. The complete Medline search syntax is included in online supplementary appendix A.

We also screened the reference lists of relevant systematic reviews conducted previously.

Grey literature was searched through the websites of 10 major humanitarian agencies and organisations who are actively involved in responding to or researching conflict situations: Emergency Nutrition Network, International Committee of the Red Cross, International Rescue Committee, Médecins Sans Frontières, Save the Children, United Nations Population Fund, United Nations High Commissioner for Refugees, UNICEF, Women's Refugee Commission and World Vision. We used broad terms for MHPSS interventions in conflict-affected populations, tailored to the search functionality of each website. Due to the large volume of grey literature available, we reviewed grey literature documents published from 1 January 2013 to 30 November 2018 only, for feasibility.

Eligibility criteria

Eligible publications were limited to those reporting on conflict-affected populations in low-income or middle-income countries, as classified by the World Bank.²³ The publication must have described an MHPSS intervention being delivered during or within 5 years of cessation of a conflict. The intervention had to target or include neonates, children, adolescents or women of reproductive age. In order to identify the most informative resources from the large volume of grey literature available, eligible grey literature publications also had to report on both the delivery site and delivery personnel for each intervention.

Non-English publications; case reports of single patients; or publications reporting on military personnel, refugee populations bound for high-income countries, surgical techniques, economic or mathematical modelling were excluded. Other excluded publications included systematic reviews, guidelines and studies where no specific health intervention was described.

Data extraction and analysis

All identified indexed records were downloaded into EndNote software and duplicates were removed. Unique records were then imported into Covidence software for screening. Titles and abstracts were reviewed in duplicate and the full-text of potentially relevant publications studies was screened for eligibility by a single reviewer.

We extracted data and information from all indexed and grey literature publications that met the eligibility criteria in duplicate, using a customised form in REDCap software. Key variables with regard to setting, target population, study design and intervention descriptions were extracted. The double-entered data were compared and any inconsistencies were resolved through discussion or by a third reviewer if needed.

Descriptive statistics were used to summarise key characteristics of the conflict and intervention, including displacement status and delivery characteristics, and we narratively synthesised information on the factors impeding or facilitating intervention delivery. We




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of literature selection.

tabulated reported intervention coverage and effectiveness data. Due to the substantial methodological heterogeneity among publications, meta-analyses were not attempted.

RESULTS

The initial electronic search yielded 37 854 unique records, with most being excluded after title and abstract screening as they did not meet the review eligibility criteria (figure 1). Of the 643 potentially relevant publications screened in full text, 125 were identified as eligible for this review. The grey literature search identified another 25 eligible publications, and a further seven were identified through reference list of previous reviews, with a total of 157 publications finally included in the review.^{24–154} Table 1 summarises the characteristics of all included publications. Further details are presented in the online supplementary appendix B.

The majority of publications reported on observational studies (n=57, 36%) or randomised controlled trials (n=29, 18%), or were non-research reports (n=53, 34%). Almost half, 43%, of the included publications (n=68) described interventions targeted at children and adolescents, while only 17% were targeted at women of reproduction age (n=26). Only six publications in the discussed interventions that were delivered to children and their caregivers simultaneously. The most common displacement status reported on was refugee populations and IDPs, and camps were the most prevalent setting of displaced populations.

A variety of mental health conditions was reported, ranging from post-traumatic stress disorder and trauma-related disorders to general mental health and

Table 1 Summary of characteristics of included studies (n=157)

Geographic region *†	n
Sub-Saharan Africa	65
Middle East and North Africa	36
Europe and Central Asia	26
South Asia	20
East Asia and Pacific	10
Latin America and the Caribbean	2
Publication type	n
Observational study	57
Non-research report	53
Randomised controlled trial	29
Non-randomised controlled trial	10
Qualitative study	7
Mixed methods	1
Target population type†	n
All/general population	77
Adolescents (10–19 years)	59
Children (13 months–9 years)	39
All women	26
Displacement status of beneficiary population†	n
Refugees	63
IDPs	56
Unreported	47
Non-displaced	16
Host	5
Setting of displaced populations	n
Unreported	69
Camp	45
Mixed	22
Dispersed	21

*World Bank Regions.

†Individual publications may contribute to multiple categories. IDP, internally displace person.

functioning, with only one publication focusing on alcohol and substance abuse (online supplementary appendix B). Almost half of the publications, 44%, reported interventions being delivered solely through non-governmental organisation (NGO) platforms, and only 10% of publications reported interventions being delivered through NGO as well as education platforms.

The included publications were geographically diverse (figure 2A). The majority reported on conflict-affected populations in Sub-Saharan Africa, Europe and South Asia, with the highest concentration of publications coming from Uganda (22 studies). Latin America was the least represented region, with only Colombia and Ecuador being represented in one publication each.




Figure 2 (A) Geographical distribution of included publications. (B) Geographical distribution of included publications, by refugee population displacement status. (C) Geographical distribution of included publications, by internally displaced population status.




Figure 3 Publication counts by publication year and intervention start year.

Stratified by the displacement status of the populations, the geographical distributions of publications reporting on refugees and IDPs followed broadly similar patterns, but no publications focused on refugees in West Africa, and more countries featured in reports of IDPs than of refugees, especially in Sub-Saharan Africa and South-East Asia (figure 2B and C).

Most of the literature on the delivery of MHPSS interventions for women and children in conflict settings is recent, with the majority published after 2012, and with peaks in 2013 and 2016 (figure 3). When examining the years that reported interventions was first delivered, the included publications are more widely dispersed over time, with peaks in 2007 and 2012.

Psychosocial support was the most frequently reported intervention delivered to all study populations, followed by training interventions and then by screening (for referral/with intention to treat) (figure 4A). The delivery of counselling, creative arts therapy and psychoeducation interventions were also reported relatively frequently. Several types of therapy including eye movement desensitisation and reprocessing (EMDR), mind-body techniques and group interpersonal psychotherapy were reported with similar, low frequency. Interventions targeted at children and adolescents followed a different pattern, where the delivery of creative arts therapy was reported most frequently, followed by psychosocial support (figure 4B).

When interventions were mapped to the places and levels of care at which they were delivered, different

patterns are evident for those interventions delivered to all populations ($n=345$) compared with those targeted at children and adolescents ($n=95$; table 2).

Among interventions delivered to populations of all ages, psychosocial support and screening for referral were delivered broadly across all sites and at different levels of care, while EMDR was only delivered in health posts/mobile clinics. Interventions targeted to children and adolescents, however, were almost exclusively delivered in specialised centres and schools, with only counselling and psychosocial support being delivered more widely. Only two interventions were delivered to children and adolescents through an outreach approach, with screening being delivered through a web-based modality, and counselling being delivered at health posts or mobile clinics. With the exception of schools, community-based delivery approaches were lacking for interventions targeted to youth, and no interventions were delivered at the home.

Different patterns for children and adolescents compared with the general population are also evident when interventions were mapped to the personnel delivering them (table 3).

Psychiatrists and physicians delivered a wide range of interventions in the general population, but only screening for referral and group therapy was delivered by these professionals to youth specifically. NGO staff/researchers and counsellors/therapists/psychologists delivered the widest range of interventions in both populations. Nurses and community health workers did not deliver any interventions to children and adolescents, and civic/religious




Figure 4 Reporting frequency of mental health and psychosocial support interventions for all ages (top) and for children aged 5–19 (bottom). EMDR, eye movement desensitisation and reprocessing; NOS, not otherwise specified.

leaders only delivered psychosocial support. In the general population, psychosocial support was delivered by all types of personnel, with psychoeducation and cognitive-behavioural therapy (CBT) also delivered by many types. Similarly, in the 5–19 year age group, CBT was delivered by the widest range of personnel, followed by screening for referral and psychosocial support.

Relatively few publications that reported on the delivery of MHPSS interventions to women and children in conflict settings also reported on intervention coverage or effectiveness. Coverage data were available from only two publications included in our review,⁹⁶⁹⁹ focusing on counselling and EMDR in adolescents and women, respectively. Quantitative data relating to the effect of interventions were available from 17 publications.^{29 33 41 55 57 67 68 84 98 106 113 141 144 146 147 149 150} Reported outcomes included anxiety-related symptoms, depression-related symptoms, general mental health and functioning, post-traumatic stress disorder and

trauma-related disorder. Nearly all interventions for which effectiveness data were reported were targeted to children and adolescents, with a majority delivered in schools and a large proportion delivered to IDPs. Most of the publications reporting quantitative data did not report final effect estimates, and the common non-reporting of measures of dispersion or correlation coefficients prevented our post-hoc calculation of such estimates. Furthermore, the heterogeneity of the settings, interventions and reported outcomes barred us from further synthesising the retrieved data on effectiveness. However, overall, school-based interventions or interventions delivered in school settings appear to be effective in reducing anxiety-related symptoms, depression-related symptoms post-traumatic stress disorder and general mental health and functioning in school-aged children. All intervention coverage and effectiveness data extracted from included publications are presented in online supplementary appendix C.

Table 2 Delivery characteristics of interventions, by age group (all ages, and 5–19 years) and site of delivery

Interventions for all ages (n=345)	Outpatient				Outreach		Community based			
	Hospitals	Clinics	Research centres	Specialised centres	Web based	Health posts/ mobile clinics	Home	Place of worship	Schools	Community centres
Counselling										
Screening for referral										
Counselling (NOS)										
Cognitive-behavioural therapy										
Creative arts therapy										
EMDR										
Group therapy										
Group interpersonal psychotherapy										
Group therapy (NOS)										
Psychotherapy										
Narrative exposure therapy										
Play therapy										
Psychoeducation										
Psychosocial support (NOS)										
Psychotherapy (NOS)										
Technique/skills										
Mind-body techniques										
Life skills/problem-solving training										
School-based intervention (NOS)										
Other interventions										
Medication										
Sensitisation campaigns/community mobilisation										
Other										
Training										

NOS, Not Otherwise Specified.

Interventions for children 5–19 years (n=95)	Outpatient				Outreach		Community based			
	Hospitals	Clinics	Research centres	Specialised centres	Web based	Health posts/ mobile clinics	Home	Place of worship	Schools	Community centres
Counselling										
Screening for referral										
Counselling (NOS)										
Cognitive-behavioural therapy										
Creative arts therapy										
EMDR										
Group therapy										
Group interpersonal psychotherapy										
Group therapy (NOS)										
Psychotherapy										
Narrative exposure therapy										
Play therapy										
Psychoeducation										
Psychosocial support (NOS)										
Psychotherapy (NOS)										
Technique/skills										
Mind-body techniques										
Life skills/problem-solving training										
School-based intervention (NOS)										

Continued

Table 2 Continued

Interventions for children 5–19 years (n=95)	Outpatient				Outreach		Community based			
	Hospitals	Clinics	Research centres	Specialised centres	Web based	Health posts/mobile clinics	Home	Place of worship	Schools	Community centres
Other interventions										
Medication										
Sensitisation campaigns/community mobilisation										
Other										
Training										

EMDR, eye movement desensitisation and reprocessing; NOS, Not otherwise specified.

A number of key barriers to and facilitators of MHPSS intervention delivery were identified from the reviewed literature and grouped into themes, with supporting excerpts from the publications presented in table 4. Barriers to the delivery of interventions included limited access to target populations as a result of security constraints, in addition to infrastructure impediments such as limited network coverage and bombing of clinics. Language and cultural barriers also existed where implementers had difficulty communicating with the target population, and vice versa. In order to manage this, some programme trained lay staff from like communities; this strategy encountered other challenges however, where community members were dealing with their own trauma, or were lacking basic education, and were therefore unable to deliver services. Other barriers relating to the heterogeneity of study populations made it difficult to implement uniform interventions to wide groups of individuals. Many tactics to overcome the reported barriers were outlined in the literature. While training of lay staff encompassed some hindrances, generally, the utilisation of local community members was an important facilitating aspect of the delivery of MHPSS interventions, where doing so allowed for increased access to and acceptability among target populations. Training of outreach volunteers was also found to be imperative for being able to reach and refer people in need to the correct services and facilities. Another facilitator in the delivery of MHPSS interventions was integrating such interventions into existing networks and adapting them to the specific context. Such integration was especially helpful when they were incorporated into school-based programme, as they created safe environments for affected youth.

DISCUSSION

This review provides a comprehensive overview of reported strategies and approaches used to deliver MHPSS interventions to conflict-affected women and children in low-income and middle-income countries, drawing on 157 indexed and grey literature publications identified using rigorous systematic review methods. Evidence syntheses in the past have examined the prevalence of mental health disorders in conflict-affected

populations^{5 8 13–19}; however, this review is the first, to our knowledge, to assess the mechanisms of intervention delivery and implementation in the particular context of women and children impacted by conflict.

While the body of literature on MHPSS in humanitarian contexts is growing, much work remains to improve the breadth of evidence on intervention effectiveness as well as effective delivery strategies to increase the coverage of interventions. The majority of the publications included in the present review reported on observational rather than experimental research, and those research studies that employed experimental designs were typically conducted in stable settings, such as well-established refugee camps, with limited generalisability to other conflict settings.

Psychosocial support, training, screening (for referral/with intent to treat) and counselling accounted for more than half of the delivered interventions reported in the included publications. Only one publication was identified that focused on interventions related to alcohol or substance abuse disorders.⁸⁷ Understanding that conflict-affected populations are at a higher risk of experiencing alcohol and drug misuse as a result of coping mechanisms,^{19 155} it is surprisingly to observe an absence of strategies to prevent substance abuse.

While more than 40% of included publications reported on interventions targeted to children and adolescents, very few reported on interventions engaging parents/caregivers along with their children. Considering the significant toll that conflict and displacement take on children and families by destroying pre-existing community structures and social networks, it is important to use parents and caregivers as natural support systems for children, rather than relying on external aid such as counsellors and teachers.¹⁵⁶ Similarly, very few studies described interventions that were promoted or delivered by religious leaders or through faith-based delivery channels. Populations affected by crises tend to use faith as a coping mechanism and express a spiritual or religious identity in an effort to seek comfort.¹² Inadequate use of these networks is a missed opportunity to increase the reach of MHPSS interventions. Additionally, with the



Table 3 Delivery characteristics of interventions, by age group (all ages, and 5–19 years) and personnel

Interventions for all ages (n=345)	Counselors/therapists/psychologists					Other mental health professionals		Health workers	Social workers	NGO staff/researchers	Facilitators/trainers	CHWs	Trained civilians/volunteers	Teachers	Civic leader/religious leaders
	Physicians/psychiatrists	Psychologists	Therapists	Nurses	Other mental health professionals	Nurses	Health workers								
Counselling															
Screening for referral															
Counselling (NOS)															
Cognitive-behavioural therapy															
Creative arts therapy															
EMDR															
Group therapy															
Group interpersonal psychotherapy															
Group therapy (NOS)															
Psychotherapy															
Narrative exposure therapy															
Play therapy															
Psychoeducation															
Psychosocial support (NOS)															
Psychotherapy (NOS)															
Technique/skills															
Mind-body techniques															
Life skills/problem-solving training															
School-based intervention (NOS)															
Other interventions															
Medication															
Sensitisation campaigns/community mobilisation															
Other															
Training															

Interventions for children 5–19 years (n=95)	Counselors/therapists/psychologists					Other mental health professionals		Health workers	Social workers	NGO staff/researchers	Facilitators/trainers	CHWs	Trained civilians/volunteers	Teachers	Civic leader/religious leaders
	Physicians/psychiatrists	Psychologists	Therapists	Nurses	Other mental health professionals	Nurses	Health workers								
Counselling															
Screening for referral															
Counselling (NOS)															

CHWs, Community Health Workers; NGO, Non-governmental organization; NOS, Not otherwise specified.

Continued

Table 3 Continued

Interventions for children 5–19years (n=95)	Physicians/ Psychiatrists	Counsellors/ Therapists/ Psychologists	Other mental health professionals	Nurses	Health workers	Social workers	NGO staff/ Researchers	Facilitators/ Trainers	CHWs	Trained civilians/ Volunteers	Teachers	Civic leader/ Religious leaders
Cognitive behavioural therapy												
Creative arts therapy												
EMDR												
Group Therapy												
Group interpersonal psychotherapy												
Group therapy (NOS)												
Psychotherapy												
Narrative exposure therapy												
Play therapy												
Psychoeducation												
Psychosocial support (NOS)												
Psychotherapy (NOS)												
Techniques/Skills												
Mind-body techniques												
Life skills/Problem solving training												
School-based intervention (NOS)												
Other Interventions												
Medication												
Sensitisation campaigns / community mobilisation												
Other												
Training												

EMDR, eye movement desensitisation and reprocessing; NGO, non-governmental organisation; NOS, not otherwise specified.

Table 4 Reported barriers to and facilitators of intervention delivery

Barriers

Theme	Examples/excerpts
Access and security	<ul style="list-style-type: none"> ▶ “A continuous problem was the security situation, which impacted implementation of the programmes from the level of access to the target population, selection of staff and development of communication, to the identification of local support mechanisms”¹³² ▶ “There are always security risks to staff members due to the presence of landmines and unexploded ordnances, in addition to general travel restrictions imposed by peace-keeping troops that directly affect the sample selection process”⁶⁹ ▶ “Time was another major constraint during these camps as, due to security concerns, travelling was not possible after dark”⁷⁶ ▶ “Another obstacle was that the program coordinators were not allowed to travel to Iraq due to the worsened security situation. They were therefore forced to organize and supervise the program via the Internet, Skype, and by telephone”¹⁵²
Infrastructure	<ul style="list-style-type: none"> ▶ “Poor network coverage and phone charging facilities in the settlement makes community mobilisation processes difficult for the implementation team”¹⁰³ ▶ “The recent bombing and closure of the clinic resulted in a huge loss for the local communities’ leaving a gap in critical services for the health system of this war ravaged region”⁷³ ▶ “In these settings, the context is likely to change abruptly at any moment, and the research subjects who are participating out of goodwill alone may no longer be available. The sudden change in political circumstances meant that the original plan to measure postintervention outcomes at 4 months after admission to the nutrition program had to be modified”¹⁰⁶
Language and culture	<ul style="list-style-type: none"> ▶ “There was a huge language barrier as two-thirds of the patients could only communicate in Pashto. Even the Pashto speaking professionals had difficulty conducting interviews because of the dialect spoken by most IDPs”⁷⁶ ▶ “Throughout the intervention programme, it was observed that the majority of the children who participated in the study could not name or explain their emotions”¹⁴⁹
Training lay staff	<ul style="list-style-type: none"> ▶ “A limitation involves training teachers who had no mental health background and whose own mental state following the war trauma was a factor we could not control for completely (despite addressing it in their training)”⁸⁴ ▶ “It was a challenge to find refugees resident in the settlement with even basic education. Many had had their schooling interrupted by the conflicts from which they fled, and most of those with some education had already left the settlement in search of employment opportunities”¹¹⁷
Heterogeneity of study population	<ul style="list-style-type: none"> ▶ “There was a broad age range, not all participants had been abducted and there was a broad range of reported psychopathology at pre-test assessment”¹¹³ ▶ “Other limitations were: not investigating the process of the intervention, not involving parents, the large size and developmental heterogeneity of the intervention groups, and the different treatment requirements for PTSD and depression”¹⁴³

Facilitators

Theme	Example/excerpts
Utilisation of local community members	<ul style="list-style-type: none"> ▶ “These psychosocial outreach volunteers represent the foundation of the mental health and psychosocial support programme. This is, in part, due to the fact that Syrian and refugee outreach volunteers are able to access areas that UNHCR staff are currently not permitted to visit, including collective shelters and communities hosting displaced Syrians. The outreach volunteers and the mobile MHPSS case managers are therefore able to provide a crucial outreach function by helping to decentralise psychosocial support and to bring mental health services closer to vulnerable populations”⁷⁰ ▶ “The use of local volunteers and program graduates promotes community as well as enables the program more flexibility to reach out to a wider range of war-affected children in Kosovo”⁶⁶ ▶ “The intervention which can be easily delivered by trained community volunteers (CVs). The CVs in our study by virtue of residing in these camps had the advantage of sharing the same language and culture of the refugees, thereby enhancing their acceptability”¹⁵¹
Training of outreach volunteers for referrals	<ul style="list-style-type: none"> ▶ “Community outreach in DRC was also an important source of referral. This outreach by MSF includes specific information on sexual violence services, which may be a reason for the high proportion of survivors of sexual violence, including children, presenting to the programme”⁹⁶ ▶ “Community-based activities allowed outreach to a wide range of people among those in need as well as those who were not in need. After defining the scope of the intervention, the coverage of the refugee population could be extended to districts other than the ones previously identified, according to the population needs”⁹¹

Continued

Table 4 Continued

Facilitators

Integration of MHPSS interventions into existing PHC/referral networks	<ul style="list-style-type: none"> ▶ “The main difference between the two projects resided in the network; in a center-based program, the creation of the referral system toward the center relied mostly on the pre-existing network of service providers (NGOs, community-based organizations, local institutions), while a community-based program created its own network, usually in collaboration with community leaders and key persons such as shopkeepers”⁹¹ ▶ “In DRC, the integrated nature of the programme and links with community health workers is reflected in the fact that almost half of all patients were referred by primary healthcare services, whereas in Iraq the focus on links to secondary and tertiary health services meant they were of more importance than primary care providers. Specific provision of information on child mental health services, by providers of those services, to groups that come into contact with children appears to improve uptake”⁹⁶
Adaptation of interventions	<ul style="list-style-type: none"> ▶ “Using therapeutic consultations, mainly at the patient’s residence and often involving the family, more than three-quarters of patients had improved at the end of therapy. The importance of this adapted therapy, based on the psychodynamic approach and respecting the Palestinians’ cultural characteristics, was possible and efficient”⁶⁵ ▶ “In the absence of trained professionals for emergency interventions, mobilizing teachers was the best alternative to reach the largest number of students”⁸⁴
School-based approach	<ul style="list-style-type: none"> ▶ “School-based intervention is the best way to reach a large number of children suffering from conflict-related distress and to target youth in their natural school environment by training professionals in the educational system. School-based programs allow youth to be trained and treated in a non-stigmatizing environment with little disruption to their daily schedules, which is important for mitigating posttraumatic stress”²⁴ ▶ “Underscoring program sustainability, the Federal Ministry incorporated implementing the program into the counselors’ job requirements and contracted local mental health professionals to provide supervision. The resulting professional network interlinked participating schools with local mental health clinics and the local university and allowed the program to run autonomously throughout the school year”⁹⁰

DRC, Democratic Republic of Congo; IDP, internally displaced person; MHPSS, mental health and psychosocial support; MSF, Médecins Sans Frontières; NGO, non-governmental organisation; PHC, primary health centre; PTSD, post-traumatic stress disorder; UNHCR, United Nations High Commissioner for Refugees.

exception of schools, other community-based delivery approaches were not widely used for interventions targeted to children and adolescents. Outreach and community-based platforms are particularly important in displacement settings as a means to strengthen and facilitate the empowerment of communities.¹⁵⁶

Almost half of the reviewed literature describe local and international NGOs working independently of each other and government bodies. While some MHPSS interventions were delivered through existing health or education systems, sustainability and increased coverage would be achieved if NGO-run programmes could be delivered through formal health and education systems and leveraging existing resources and structures,^{17 74} considering humanitarian donor funds are generally short term. According to the WHO Mental Health Gap Action Programme guidelines, integration can be done by training healthcare staff in the identification, management and appropriate referral of mental health cases.¹⁵⁷ Such integration will ensure sustainability, increase accessibility, and reduce the associated stigma of mental health. Accessibility and reach of MHPSS interventions could be further amplified by packaging them with interventions in other sectors, such as nutrition. In 2006, WHO recommended integrating psychosocial stimulation into emergency feeding programme, for example, yet not many studies have evaluated the impact of such combined interventions in emergency

settings.^{18 158} Multisectoral integration is an area that should be prioritised for MHPSS, as the most vulnerable populations tend to suffer from comorbidities and might be more easily reached and more effectively helped by targeting these simultaneously.¹⁰⁶ Moreover, this integration can help ensure programme sustainability and thereby reduce the unintended negative consequences of suspending or withdrawing MHPSS interventions that communities were previously able to access.

Additionally, given that language and culture are cited in the literature as key intervention delivery barriers, more evidence on the unintended consequences of delivering MHPSS interventions that are culturally insensitive is needed.¹⁵⁹ Cultural sensitivity needs to be prioritised in the context of diagnosing mental health disorders in conflict settings, since the Diagnostic and Statistical Manual of Mental Disorders is developed for western settings.¹⁶⁰ Cultural appropriateness can be achieved with high-quality training of implementors and mental health professionals specific to the context they are working in, and also through increased delivery of web-based or mobile app interventions where opportunities for face-to-face interactions are limited.⁶

A major limitation of the literature is the sizeable number of publications that do not report key information such as the site of intervention delivery or the personnel delivering the intervention, indicating the

need for stronger scientific reporting. There was also much heterogeneity and non-standardisation in the reporting of intervention outcomes, making pooled analysis challenging. In addition to the limitations of the literature, several methodological limitations of this systematic review must also be noted. Due to capacity and feasibility, we limited our review to English-language publications only, and our search of the grey literature was comprehensive and systematic, but not exhaustive. We may therefore have missed other publications that may have contributed different information to this review.

CONCLUSION

The growing literature, including among children, is encouraging, but more attention must be paid to establishing and documenting the research and practice of MHPSS delivery in a wider range of conflict contexts covering a wider range of interventions. Better leveraging of existing resources and structures in the health and the education sector, and others such as nutrition, could help improve the reach of MHPSS interventions and also their sustainability. So, too, could an increased focus on interventions that make better use of parents' / caregivers' supportive roles in their children's lives, as well as other pre-existing or emerging social support networks, and that rely less on the availability of specialised mental health personnel. More robust evaluation of intervention effectiveness as well as intervention delivery effectiveness would improve the evidence base, and could ultimately improve the mental health and psychosocial well-being of conflict-affected women and children globally.

Twitter Fahad J Siddiqui @fjsepi

Contributors ZAB, MFG and FJS conceptualised the review. FJS created the search strategy and conducted the literature search in electronic databases. MK, SM, DA, MM, RPJ and SS screened all studies for inclusion into the systematic review and performed data extraction. MK conducted the analysis with support from MFG and FJS. MK drafted the manuscript and MFG and ZAB critically revised it. All authors approved the final version.

Funding As coordinator of the BRANCH Consortium (Bridging Research & Action in Conflict Settings for the Health of Women & Children), the SickKids Centre for Global Child Health has received funding for BRANCH research activities, including this systematic review, from the International Development Research Centre (108416-002 and 108640-001), the Norwegian Agency for Development Cooperation (Norad) (QZA-16/0395), the Bill & Melinda Gates Foundation (OPP1171560) and UNICEF (PCA 20181204). Aga Khan University has received funding for BRANCH activities from the Family Larsson-Rosenquist Foundation.

Disclaimer Dissemination of results to study participants or patient organisations is not applicable for this review.

Map disclaimer The depiction of boundaries on the map(s) in this article does not imply the expression of any opinion whatsoever on the part of BMJ (or any member of its group) concerning the legal status of any country, territory, jurisdiction or area or of its authorities. The map(s) are provided without any warranty of any kind, either express or implied.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethics approval was not required for this systematic review of the published literature.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Reena P Jain <http://orcid.org/0000-0001-7004-5093>

REFERENCES

- 1 Save the Children's End of Childhood Report 2018. Save the Children 2019.
- 2 People affected by conflict: humanitarian needs in numbers. Belgium: centre for research on the epidemiology of disasters 2013.
- 3 Steel Z, Chey T, Silove D, *et al*. Association of torture and other potentially traumatic events with mental health outcomes among populations exposed to mass conflict and displacement. *JAMA* 2009;302:537–49.
- 4 Mushtaq R, Shah T, Mushtaq S, *et al*. Post-Traumatic stress disorder (PTSD) in children of conflict region of Kashmir (India): a review. *J Clin Diagn Res* 2016;10:VE01–3.
- 5 Turrini G, Purgato M, Ballette F, *et al*. Common mental disorders in asylum seekers and refugees: umbrella review of prevalence and intervention studies. *Int J Ment Health Syst* 2017;11:51.
- 6 Who. *Mental health promotion and mental health care in refugees and migrants: policy brief*, 2018.
- 7 Slone M, Mann S. Effects of war, terrorism and armed conflict on young children: a systematic review. *Child Psychiatry Hum Dev* 2016;47:950–65.
- 8 Shaar KH. Post-Traumatic stress disorder in adolescents in Lebanon as wars gained in ferocity: a systematic review. *J Public Health Res* 2013;2:e17.
- 9 Blanchet K, Ramesh A, Frison S, *et al*. Evidence on public health interventions in humanitarian crises. *The Lancet* 2017;390:2287–96.
- 10 Augustinavicius JL, Greene MC, Lakin DP, *et al*. Monitoring and evaluation of mental health and psychosocial support programs in humanitarian settings: a scoping review of terminology and focus. *Confl Health* 2018;12.
- 11 IASC. *Inter-Agency standing Committee (IASC) guidelines on mental health and psychosocial support in emergency setting*. Geneva: IASC, 2007.
- 12 SPHERE. *The sphere Handbook: Humanitarian charter and minimum Standards in humanitarian response*, 2018.
- 13 Attanayake V, McKay R, Joffres M, *et al*. Prevalence of mental disorders among children exposed to war: a systematic review of 7,920 children. *Med Confl Surviv* 2009;25:4–19.
- 14 Bogic M, Njoku A, Priebe S. Long-Term mental health of war-refugees: a systematic literature review. *BMC Int Health Hum Rights* 2015;15:29.
- 15 Fellmeth G, Fazel M, Plugge E. Migration and perinatal mental health in women from low- and middle-income countries: a systematic review and meta-analysis. *BJOG: Int J Obstet Gy* 2017;124:742–52.
- 16 Vossoughi N, Jackson Y, Gusler S, *et al*. Mental health outcomes for youth living in refugee camps: a review. *Trauma Violence Abuse* 2018;19:528–42.
- 17 Tol WA, Barbui C, Galappatti A, *et al*. Mental health and psychosocial support in humanitarian settings: linking practice and research. *The Lancet* 2011;378:1581–91.
- 18 Jordans MJD, Tol WA, Komproe IH, *et al*. Systematic review of evidence and treatment approaches: psychosocial and mental health care for children in war. *Child Adolesc Ment Health* 2009;14:2–14.
- 19 Horyniak D, Melo JS, Farrell RM, *et al*. Epidemiology of substance use among forced migrants: a global systematic review. *PLoS One* 2016;11:e0159134.
- 20 Charlson F, van Ommeren M, Flaxman A, *et al*. New who prevalence estimates of mental disorders in conflict settings: a systematic review and meta-analysis. *The Lancet* 2019;394:240–8.
- 21 Jordans MJD, Pigott H, Tol WA. Interventions for children affected by armed conflict: a systematic review of mental health and

- psychosocial support in low- and middle-income countries. *Curr Psychiatry Rep* 2016;18.
- 22 Tol WA, Stavrou V, Greene MC, *et al*. Sexual and gender-based violence in areas of armed conflict: a systematic review of mental health and psychosocial support interventions. *Confl Health* 2013;7.
 - 23 Bank W. Income classification, 2018. Available: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
 - 24 Abdeen Z, Qasrawi R, Nabil S, *et al*. Psychological reactions to Israeli occupation: findings from the national study of school-based screening in Palestine. *Int J Behav Dev* 2008;32:290–7.
 - 25 Abramowitz SA. The poor have become rich, and the rich have become poor: collective trauma in the Guinean Languette. *Soc Sci Med* 2005;61:2106–18.
 - 26 Acarturk C, Konuk E, Cetinkaya M, *et al*. EMDR for Syrian refugees with posttraumatic stress disorder symptoms: results of a pilot randomized controlled trial. *Eur J Psychotraumatol* 2015;6:27414.
 - 27 Acarturk C, Konuk E, Cetinkaya M, *et al*. The efficacy of eye movement desensitization and reprocessing for post-traumatic stress disorder and depression among Syrian refugees: results of a randomized controlled trial. *Psychol Med* 2016;46:2583–93.
 - 28 Ager A, Akesson B, Stark L, *et al*. The impact of the school-based psychosocial structured activities (PSSA) program on conflict-affected children in northern Uganda. *J Child Psychol Psychiatry* 2011;52:1124–33.
 - 29 Annan J, Sim A, Puffer ES, *et al*. Improving mental health outcomes of Burmese migrant and displaced children in Thailand: a community-based randomized controlled trial of a parenting and family skills intervention. *Prevention Science* 2017;18:793–803.
 - 30 Awodola B. An examination of methods to reintegrate former child soldiers in Liberia. *Intervention* 2012;10:30–42.
 - 31 Ayoughi S, Missmahl I, Weierstall R, *et al*. Provision of mental health services in resource-poor settings: a randomised trial comparing counselling with routine medical treatment in North Afghanistan (Mazar-e-Sharif). *BMC Psychiatry* 2012;12.
 - 32 Badri A, Crutzen R, Van den Borne HW. Exposures to war-related traumatic events and post-traumatic stress disorder symptoms among displaced Darfuri female university students: an exploratory study. *BMC Public Health* 2012;12:603.
 - 33 Bass JK, Annan J, Mclvor Murray S, *et al*. Controlled trial of psychotherapy for Congolese survivors of sexual violence. *N Engl J Med* 2013;368:2182–91.
 - 34 Bass J, Poudyal B, Tol W, *et al*. A controlled trial of problem-solving counseling for war-affected adults in Aceh, Indonesia. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:279–91.
 - 35 Bastin P, Bastard M, Rossel L, *et al*. Description and predictive factors of individual outcomes in a refugee cAMP based mental health intervention (Beirut, Lebanon). *PLoS One* 2013;8:e54107.
 - 36 Behrens GA. Use of traditional and nontraditional instruments with traumatized children in Bethlehem, West bank. *Music Therapy Perspectives* 2012;30:196–202.
 - 37 Benjamin JA. Aids prevention for refugees. The case of Rwandans in Tanzania. *Aids captions* 1996;3:4–9.
 - 38 Betancourt TS, Borisova II, Williams TP, *et al*. Sierra Leone's Former Child Soldiers: A Follow-Up Study of Psychosocial Adjustment and Community Reintegration. *Child Dev* 2010;81:1077–95.
 - 39 Betancourt TS, Newnham EA, Brennan RT, *et al*. Moderators of treatment effectiveness for war-affected youth with depression in northern Uganda. *J Adolesc Health* 2012;51:544–50.
 - 40 Betancourt TS, Newnham EA, McBain R, *et al*. Post-Traumatic stress symptoms among former child soldiers in Sierra Leone: follow-up study. *Br J Psychiatry* 2013;203:196–202.
 - 41 Betancourt TS, Yudron M, Wheaton W, *et al*. Caregiver and adolescent mental health in Ethiopian Kunama refugees participating in an emergency education program. *J Adolesc Health* 2012;51:357–65.
 - 42 Bhardwaj A, Bourey C, Rai S, *et al*. Interpersonal violence and suicidality among former child soldiers and war-exposed civilian children in Nepal. *Glob Ment Health* 2018;5:e9.
 - 43 Bolton P, Lee C, Haroz EE, *et al*. A transdiagnostic community-based mental health treatment for comorbid disorders: development and outcomes of a randomized controlled trial among Burmese refugees in Thailand. *PLoS Med* 2014;11:e1001757-e.
 - 44 Chase R, Bush K. The mental health of war affected children: a community based rehabilitation and reconciliation programme in Sri Lanka's eastern Province. *Asia Pacific Disability* 2002;50–66.
 - 45 Chauvin L, Mugaju J, Comlavi J. Evaluation of the psychosocial trauma recovery programme in Rwanda. *Eval Program Plann* 1998;21:385–92.
 - 46 Ciaessens LF, de Graaff DC, Jordans MJD, *et al*. Participatory evaluation of psychosocial interventions for children: a pilot study in northern Uganda. *Intervention* 2012;10:43–58.
 - 47 Cliff J, Noormahomed AR. The impact of war on children's health in Mozambique. *Soc Sci Med* 1993;36:843–8.
 - 48 Cohen RA, Agosin A. Common threads: a recovery programme for survivors of gender based violence. *Intervention: International Journal of Mental Health, Psychosocial Work & Counselling in Areas of Armed Conflict* 2013;11:157–68.
 - 49 Cohen S, Asgary R. Community coping strategies in response to hardship and human rights abuses among Burmese refugees and migrants at the Thai-Burmese border: a qualitative approach. *Fam Community Health* 2016;39:75–81.
 - 50 Constandinides D, Kamens S, Marshoud B, *et al*. Research in ongoing conflict zones: effects of a school-based intervention for Palestinian children. *Peace and Conflict: Journal of Peace Psychology* 2011;17:270–302.
 - 51 Corbin J, Miller J. Collaborative psychosocial capacity building in northern Uganda. *Families in Society* 2009;90:103–9.
 - 52 Cox J, Davies DR, Burlingame GM, *et al*. Effectiveness of a Trauma/Grief-Focused group intervention: a qualitative study with War-Exposed Bosnian adolescents. *Int J Group Psychother* 2007;57:319–45.
 - 53 Dahl S, Mutapcic A, Schei B. Traumatic events and predictive factors for posttraumatic symptoms in displaced Bosnian women in a war zone. *J Trauma Stress* 1998;11:137–45.
 - 54 de Jong JP, Scholte WF, Koeter MWJ, *et al*. The prevalence of mental health problems in Rwandan and Burundese refugee camps. *Acta Psychiatr Scand* 2000;102:171–7.
 - 55 Diab M, Peltonen K, Qouta SR, *et al*. Effectiveness of psychosocial intervention enhancing resilience among war-affected children and the Moderating role of family factors. *Child Abuse Negl* 2015;40:24–35.
 - 56 Donev D, Onceva S, Gligorov I. Refugee crisis in Macedonia during the Kosovo conflict in 1999. *Croat Med J* 2002;43:184–9.
 - 57 Dybdahl R. Children and mothers in war: an outcome study of a psychosocial intervention program. *Child Dev* 2001;72:1214–30.
 - 58 Dybdahl R, Pasagic I. Traumatic experiences and psychological reactions among women in Bosnia during the war. *Med Confl Surviv* 2000;16:281–90.
 - 59 Eiling E, Van Diggele-Holtland M, Van Yperen T, *et al*. Psychosocial support for children in the Republic of South Sudan: an evaluation outcome. *Intervention* 2014;12:61–75.
 - 60 Elhabiby MM, Radwan DN, Okasha TA, *et al*. Psychiatric disorders among a sample of internally displaced persons in South Darfur. *Int J Soc Psychiatry* 2015;61:358–62.
 - 61 Ertl V, Pfeiffer A, Schauer E, *et al*. Community-implemented trauma therapy for former child soldiers in northern Uganda: a randomized controlled trial. *JAMA* 2011;306:503–12.
 - 62 Espié E, Gaboulaud V, Baubet T, *et al*. Trauma-Related psychological disorders among Palestinian children and adults in Gaza and West bank, 2005-2008. *Int J Ment Health Syst* 2009;3:21.
 - 63 Firdosi M, Margoob M. Socio-Demographic profile and psychiatric comorbidity in patients with a diagnosis of post traumatic stress disorder – a study from Kashmir Valley. *Acta Medica International* 2016;3:97–100.
 - 64 Frijak A, Cengic S, Hauser M, *et al*. Gynecological complaints and war traumas. A study from Zenica, Bosnia-Herzegovina during the war. *Acta Obstet Gynecol Scand* 1997;76:350–4.
 - 65 Gaboulaud V, Reynaud C, Moro M-R, *et al*. Psychological support for Palestinian children and adults: an analysis of data from people referred to the Médecins sans Frontières programme for behavioural and emotional disorders in the occupied Palestinian Territory. *Intervention* 2010;8:131–42.
 - 66 Gerber MM, Hogan LR, Maxwell K, *et al*. Children after war: a novel approach to promoting resilience through music. *Traumatology* 2014;20:112–8.
 - 67 Gordon JS, Staples JK, Blyta A, *et al*. Treatment of posttraumatic stress disorder in postwar Kosovo high school students using Mind-body skills groups: a pilot study. *J Trauma Stress* 2004;17:143–7.
 - 68 Gordon JS, Staples JK, Blyta A, *et al*. Treatment of posttraumatic stress disorder in postwar Kosovar adolescents using Mind-body skills groups. *J Clin Psychiatry* 2008;69:1469–76.
 - 69 Gupta L, Zimmer C. Psychosocial intervention for war-affected children in Sierra Leone. *Br J Psychiatry* 2008;192:212–6.
 - 70 Harrison S, Dahman R, Ismail M, *et al*. 'Against all odds': UNHCR's mental health and psychosocial support programme for Iraqi refugees and internally displaced Syrians. *Intervention: International Journal of Mental Health, Psychosocial Work & Counselling in Areas of Armed Conflict* 2013;11:190–4.

- 71 Hasanovic M, Sinanovic O, Pajevic I, *et al*. Post-War mental health promotion in Bosnia-Herzegovina. *Psychiatry Danubina* 2006;18:74–8.
- 72 Hashemi B, Ali S, Awaad R, *et al*. Facilitating mental health screening of war-torn populations using mobile applications. *Soc Psychiatry Psychiatr Epidemiol* 2017;52:27–33.
- 73 Hemat H, Shah S, Isaakidis P, *et al*. Before the bombing: high burden of traumatic injuries in Kunduz trauma center, Kunduz, Afghanistan. *PLoS One* 2017;12:e0165270.
- 74 Hijazi Z, Weissbecker I, Chammy R. The integration of mental health into primary health care in Lebanon. *Intervention* 2011;9:265–78.
- 75 Hrnjica S, Powell SD-BE. Programs of psychosocial assistance for children with special needs and their parents after the war. *The psychosocial consequences of war: Results of empirical research from the territory of former Yugoslavia* 2000:281–3.
- 76 Humayun A, Azad N, Haq I, *et al*. Mental health and psychosocial support for the internally displaced persons in Bannu, Pakistan. *Intervention* 2016;14:33–49.
- 77 Hustache S, Moro M-R, Roptin J, *et al*. Evaluation of psychological support for victims of sexual violence in a conflict setting: results from Brazzaville, Congo. *Int J Ment Health Syst* 2009;3:7.
- 78 Jan M, Rather Y, Majeed N, *et al*. Psychosocial risk factors and clinical profile associated with attempted suicide in young adult and adolescent patients in conflict zone-Kashmir. *Annals of Tropical Medicine and Public Health* 2017;10:154–9.
- 79 Jandric S. Injury severity and functional outcome following paediatric trauma in war conditions. *Pediatr Rehabil* 2001;4:169–75.
- 80 Jafee-Bahloul H, Butler J. Use of telepsychiatry in areas of conflict: the Syrian refugee crisis as an example. *J Telemed Telecare* 2014;20:167–8.
- 81 Jones L, Rrustemi A, Shahini M, *et al*. Mental health services for war-affected children: report of a survey in Kosovo. *Br J Psychiatry* 2003;183:540–6.
- 82 Jordans MJD, Komproe IH, Tol WA, *et al*. Evaluation of a classroom-based psychoeducation intervention in conflict-affected Nepal: a cluster randomized controlled trial. *J Child Psychol Psychiatry* 2010;51:818–26.
- 83 Jordans MJD, Tol WA, Ndayisaba A, *et al*. A controlled evaluation of a brief parenting psychoeducation intervention in Burundi. *Soc Psychiatry Psychiatr Epidemiol* 2013;48:1851–9.
- 84 Karam EG, Fayyad J, Karam AN, *et al*. Effectiveness and specificity of a classroom-based group intervention in children and adolescents exposed to war in Lebanon. *World Psychiatry* 2008;7:103–9.
- 85 Khan MN, Dherani M, Chiumento A, *et al*. Evaluating feasibility and acceptability of a local psycho-educational intervention for pregnant women with common mental problems affected by armed conflict in Swat, Pakistan: a parallel randomized controlled feasibility trial. *Int J Soc Psychiatry* 2017;63:724–35.
- 86 Kohrt BA, Jordans MJD, Tol WA, *et al*. Validation of cross-cultural child mental health and psychosocial research instruments: adapting the depression self-rating scale and child PTSD symptom scale in Nepal. *BMC Psychiatry* 2011;11:127.
- 87 Lai L. Treating substance abuse as a consequence of conflict and displacement: a call for a more inclusive global mental health. *Medicine, Conflict and Survival* 2014;30:182–9.
- 88 Lama S, François K, Marwan Z, *et al*. Impact of the Syrian crisis on the hospitalization of Syrians in a psychiatric setting. *Community Ment Health J* 2016;52:84–93.
- 89 Layne CM, Pynoos RS, Saltzman WR, *et al*. Trauma/grief-focused group psychotherapy: school-based postwar intervention with traumatized Bosnian adolescents. *Group Dynamics: Theory, Research, and Practice* 2001;5:277–90.
- 90 Layne CM, Saltzman WR, Poppleton L, *et al*. Effectiveness of a school-based group psychotherapy program for war-exposed adolescents: a randomized controlled trial. *J Am Acad Child Adolesc Psychiatry* 2008;47:1048–62.
- 91 Le Roch K, Pons E, Squire J, *et al*. Two psychosocial assistance approaches for Iraqi urban refugees in Jordan and Lebanon: center-based services compared to community outreach services. *J Muslim Ment Health* 2010;5:99–119.
- 92 Lemaire J-M, Despret V, Boszormenyi-Nagy B-NC, *et al*. Residual resources, and an extensive context of trust: creating a network in a refugee cAMP in former Yugoslavia. *Int J Ment Health Nurs* 2001;30:22–6.
- 93 Liddell BJ, Silove D, Tay K, *et al*. Achieving convergence between a community-based measure of explosive anger and a clinical interview for intermittent explosive disorder in Timor-Leste. *J Affect Disord* 2013;150:1242–6.
- 94 Liebling H, Davidson L, Akello GF, *et al*. The experiences of survivors and trauma counselling service providers in northern Uganda: implications for mental health policy and legislation. *Int J Law Psychiatry* 2016;49:84–92.
- 95 Llosa AE, Casas G, Thomas H, *et al*. Short and longer-term psychological consequences of operation cast lead: documentation from a mental health program in the Gaza strip. *Confl Health* 2012;6:8.
- 96 Lokuge K, Shah T, Pintaldi G, *et al*. Mental health services for children exposed to armed conflict: Médecins Sans Frontières' experience in the Democratic Republic of Congo, Iraq and the occupied Palestinian territory. *Paediatr Int Child Health* 2013;33:259–72.
- 97 Lorschiedter A, Canavera CCW. Community based volunteers as partners for agencies working with formerly abducted children and youth: experiences from Northern Uganda. *Intervention* 2007;5:244–9.
- 98 Loughry M, Ager A, Flouri E, *et al*. The impact of structured activities among Palestinian children in a time of conflict. *J Child Psychol Psychiatry* 2006;47:060907041921002-???
- 99 Mankuta D, Aziz-Suleyman A, Yochai L, *et al*. Field evaluation and treatment of short-term psycho-medical trauma after sexual assault in the Democratic Republic of Congo. *Isr Med Assoc J* 2012;14:653–7.
- 100 McKell C, Hankir A, Abu-Zayed I, *et al*. Barriers to accessing and consuming mental health services for Palestinians with psychological problems residing in refugee camps in Jordan. *Psychiatry Danubina* 2017;29:157–63.
- 101 Mercer SW, Ager A, Ruwanpura E. Psychosocial distress of Tibetans in exile: integrating Western interventions with traditional beliefs and practice. *Soc Sci Med* 2005;60:179–89.
- 102 Mirghani Z A. Healing through sharing: an outreach project with Iraqi refugee volunteers in Syria. *Intervention: International Journal of Mental Health, Psychosocial Work & Counselling in Areas of Armed Conflict* 2013;11:321–9.
- 103 Mogga R. Addressing gender based violence and psychosocial support among South Sudanese refugee settlements in northern Uganda. *Intervention* 2017;15:9–16.
- 104 Mooren TTM, de Jong K, Kleber RJ, *et al*. The efficacy of a mental health program in Bosnia-Herzegovina: impact on coping and general health. *J Clin Psychol* 2003;59:57–69.
- 105 Morof DF, Sami S, Mangeni M, *et al*. A cross-sectional survey on gender-based violence and mental health among female urban refugees and asylum seekers in Kampala, Uganda. *Int J Gynaecol Obstet* 2014;127:138–43.
- 106 Morris J, Jones L, Berrino A, *et al*. Does combining infant stimulation with emergency feeding improve psychosocial outcomes for displaced mothers and babies? A controlled evaluation from Northern Uganda. *Am J Orthopsychiatry* 2012;82:349–57.
- 107 Mueller Y, Cristofani S, Rodriguez C, *et al*. Integrating mental health into primary care for displaced populations: the experience of Mindanao, Philippines. *Confl Health* 2011;5.
- 108 Nakimuli-Mpungu E, Alderman S, Kinyanda E, *et al*. Implementation and scale-up of Psycho-Trauma centers in a Post-Conflict area: a case study of a Private–Public partnership in northern Uganda. *PLoS Med* 2013;10:e1001427-e.
- 109 Neuner F, Onyut PL, Ertl V, *et al*. Treatment of posttraumatic stress disorder by trained lay counselors in an African refugee settlement: a randomized controlled trial. *J Consult Clin Psychol* 2008;76:686–94.
- 110 Neuner F, Schauer M, Klaschik C, *et al*. A comparison of narrative exposure therapy, supportive counseling, and psychoeducation for treating posttraumatic stress disorder in an African refugee settlement. *J Consult Clin Psychol* 2004;72:579–87.
- 111 Neuner F, Schauer M, Roth WT, *et al*. A narrative exposure treatment as intervention in a refugee cAMP: a case report. *Behav Cogn Psychother* 2002;30:205–9.
- 112 Newnham EA, McBain RK, Hann K, *et al*. The youth readiness intervention for war-affected youth. *J Adolesc Health* 2015;56:606–11.
- 113 O'Callaghan P, Branham L, Shannon C, *et al*. A pilot study of a family focused, psychosocial intervention with war-exposed youth at risk of attack and abduction in north-eastern Democratic Republic of Congo. *Child Abuse Negl* 2014;38:1197–207.
- 114 O'Callaghan P, McMullen J, Shannon C, *et al*. Comparing a trauma focused and non trauma focused intervention with war affected Congolese youth. *Intervention* 2015;13:28–44.
- 115 O'Brien DP, Mills C, Hamel C, *et al*. Universal access: the benefits and challenges in bringing integrated HIV care to isolated and conflict affected populations in the Republic of Congo. *Confl Health* 2009;3:1–10.

- 116 Okello J, Onen TS, Musisi S. Psychiatric disorders among war-abducted and non-abducted adolescents in Gulu district, Uganda: a comparative study. *Afr J Psychiatry* 2007;10:225–31.
- 117 Onyut LP, Neuner F, Schauer E, *et al*. The Nakivale cAMP mental health project: building local competency for psychological assistance to traumatised refugees. *Intervention: International Journal of Mental Health, Psychosocial Work & Counselling in Areas of Armed Conflict* 2004;2:90–107.
- 118 Ovuga E, Oyok TO, Moro EB. Post traumatic stress disorder among former child soldiers attending a rehabilitative service and primary school education in northern Uganda. *Afr Health Sci* 2008;8:136–41.
- 119 Panter-Brick C, Dajani R, Eggerman M, *et al*. Insecurity, distress and mental health: experimental and randomized controlled trials of a psychosocial intervention for youth affected by the Syrian crisis. *Journal of Child Psychology and Psychiatry* 2018;59:523–41.
- 120 Peltonen K, Qouta S, El Sarraj E, *et al*. Effectiveness of school-based intervention in enhancing mental health and social functioning among war-affected children. *Traumatology* 2012;18:37–46.
- 121 Qouta SR, Palosaari E, Diab M, *et al*. Intervention effectiveness among war-affected children: a cluster randomized controlled trial on improving mental health. *J Trauma Stress* 2012;25:288–98.
- 122 Quosh C. Mental health, forced displacement and recovery: integrated mental health and psychosocial support for urban refugees in Syria. *Intervention* 2013;11:295–320.
- 123 Rabaia Y, Giacaman R, Nguyen-Gillham V. Violence and adolescent mental health in the occupied Palestinian Territory: a contextual approach. *Asia Pac J Public Health* 2010;22:216S–21.
- 124 Rasmussen A, Nguyen L, Wilkinson J, *et al*. Rates and impact of trauma and current stressors among Darfuri refugees in eastern Chad. *Am J Orthopsychiatry* 2010;80:227–36.
- 125 Rees B, Travis F, Shapiro D, *et al*. Significant Reductions in Posttraumatic Stress Symptoms in Congolese Refugees Within 10 days of Transcendental Meditation Practice. *J Trauma Stress* 2014;27:112–5.
- 126 Reiffers R, Dahal RP, Koirala S, *et al*. Psychosocial support for Bhutanese refugees in Nepal. *Intervention* 2013;11:169–79.
- 127 Roberts B, Makhshvili N, Javakhishvili J, *et al*. Mental health care utilisation among internally displaced persons in Ukraine: results from a nation-wide survey. *Epidemiol Psychiatr Sci* 2017;1:1–12.
- 128 Roberts B, Ocaka KF, Browne J, *et al*. Factors associated with post-traumatic stress disorder and depression amongst internally displaced persons in northern Uganda. *BMC Psychiatry* 2008;8:38.
- 129 Salah TTM, Abdelrahman A, Lien L, *et al*. The mental health of internally displaced persons: an epidemiological study of adults in two settlements in central Sudan. *International Journal of Social Psychiatry* 2013;59:782–8.
- 130 Salem-Pickartz J. Peer counsellors training with refugees from Iraq: a Jordanian case study. *Intervention* 2007;5:232–43.
- 131 Schei B, Dahl S, Dahl WY. The burden left my heart: psycho-social services among refugee women in Zenica and Tuzla, Bosnia-Herzegovina during the war. *Women & Therapy* 1999;22:139–51.
- 132 Scholte WF, van de Put W, de Jong JP, *et al*. A protocol for psychosocial intervention in refugee crisis: early experiences in Rwandan refugee camps. *Intervention: International Journal of Mental Health, Psychosocial Work & Counselling in Areas of Armed Conflict* 2004;2:181–92.
- 133 Sehovic M, Powell SD-BE. *Evaluation of results of cognitive-behavioural therapy with traumatised children of displaced persons*. The psychosocial consequences of war: Results of empirical research from the territory of former Yugoslavia, 2000: 263–6.
- 134 Shah S, Van den Bergh R, Van Bellinghen B, *et al*. Offering mental health services in a conflict affected region of Pakistan: who comes, and why? *PLoS One* 2014;9:e97939.
- 135 Shaheen MMA, Oppenheim S, Abdeen AB-T. Youth resilience makes a difference in mitigating stress: teacher mediated school intervention in Bethlehem. *Intervention: Journal of Mental Health and Psychosocial Support in Conflict Affected Areas* 2016;14:305–19.
- 136 Sheikh TL, Mohammed A, Agunbiade S, *et al*. Psycho-trauma, psychosocial adjustment, and symptomatic post-traumatic stress disorder among internally displaced persons in Kaduna, northwestern Nigeria. *Front Psychiatry* 2014;5:127.
- 137 Sliep Y, Weingarten K, Gilbert A. Narrative theatre as an interactive community approach to mobilizing collective action in northern Uganda. *Families, Systems, & Health* 2004;22:306–20.
- 138 Sonderegger R, Rombouts S, Ocen B, *et al*. Trauma rehabilitation for war-affected persons in northern Uganda: a pilot evaluation of the empower programme. *Br J Clin Psychol* 2011;50:234–49.
- 139 Stuvland R, Durakovic-Belko E, Powell SD-BE. Evaluation of UNICEF-supported school psychosocial programs in B&H (Bosnia and Herzegovina) 1993–1999. *The psychosocial consequences of war: Results of empirical research from the territory of former Yugoslavia* 2000:234–7.
- 140 Szilard I, Cserti A, Hoxha R, *et al*. International organization for migration: experience on the need for medical evacuation of refugees during the Kosovo crisis in 1999. *Croat Med J* 2002;43:195–8.
- 141 Tanaka A, Alaimo A-S. Assessment of the psychosocial development of children attending nursery schools in Karen refugee camps in Thailand. *International Journal of Early Childhood* 2013;45:279–305.
- 142 Tayler-Smith K, Zachariah R, Hinderaker SG, *et al*. Sexual violence in post-conflict Liberia: survivors and their care. *Tropical Medicine & International Health* 2012;17:1356–60.
- 143 Thabet AA, Vostanis P, Karim K. Group crisis intervention for children during ongoing war conflict. *Eur Child Adolesc Psychiatry* 2005;14:262–9.
- 144 Tol WA, Komproue IH, Jordans MJD, *et al*. School-Based mental health intervention for children in war-affected Burundi: a cluster randomized trial. *BMC Med* 2014;12:56.
- 145 Tol WA, Komproue IH, Jordans MJD, *et al*. Brief multi-disciplinary treatment for torture survivors in Nepal: a naturalistic comparative study. *International Journal of Social Psychiatry* 2009;55:39–56.
- 146 Tol WA, Komproue IH, Jordans MJD, *et al*. Outcomes and moderators of a preventive schoolbased mental health intervention for children affected by war in Sri Lanka: a cluster randomized trial. *World Psychiatry* 2012;11:114–22.
- 147 Tol WA, Komproue IH, Susanty D, *et al*. School-Based mental health intervention for children affected by political violence in Indonesia. *JAMA* 2008;300:655–62.
- 148 Tufan AE, Alkin M, Bosgelmez S. Post-Traumatic stress disorder among asylum seekers and refugees in Istanbul may be predicted by torture and loss due to violence. *Nord J Psychiatry* 2013;67:219–24.
- 149 Ugurlu N, Akca L, Acarturk C, *et al*. An art therapy intervention for symptoms of post-traumatic stress, depression and anxiety among Syrian refugee children. *Vulnerable Child Youth Stud* 2016;11:89–102.
- 150 Veronese G, Barola G. Healing stories: an expressive-narrative intervention for strengthening resilience and survival skills in school-aged child victims of war and political violence in the Gaza strip. *Clin Child Psychol Psychiatry* 2018;23:311–32.
- 151 Vijayakumar L, Mohanraj R, Kumar S, *et al*. CASP - An intervention by community volunteers to reduce suicidal behaviour among refugees. *Int J Soc Psychiatry* 2017;63:589–97.
- 152 Wagner B, Schulz W, Knaevelsrud C. Efficacy of an Internet-based intervention for posttraumatic stress disorder in Iraq: a pilot study. *Psychiatry Res* 2012;195:85–8.
- 153 Wessells M, Monteiro C, Ahearn A-M-S. Psychosocial assistance for youth: toward reconstruction for peace in Angola. *J Soc Issues* 2006;62:121–39.
- 154 Yeomans PD, Forman EM, Herbert JD, *et al*. A randomized trial of a reconciliation workshop with and without PTSD psychoeducation in Burundian sample. *J Trauma Stress* 2010;9:n/a–12.
- 155 Ezard N. Substance use among populations displaced by conflict: a literature review. *Disasters* 2012;36:533–57.
- 156 UNHCR. *Community-based Protection and Mental Health & Psychosocial Support*. UNHCR, 2017.
- 157 WHO. *Mental health gap action programme (mhGAP): operations manual*. Geneva, 2018.
- 158 WHO. *Mental health and psychosocial Well-Being among children in severe food shortage situations*. Geneva, 2006.
- 159 Lie DA, Lee-Rey E, Gomez A, *et al*. Does cultural competency training of health professionals improve patient outcomes? A systematic review and proposed algorithm for future research. *J Gen Intern Med* 2011;26:317–25.
- 160 Kupfer D, First M, Regier D. *A research agenda for DSM-V*. Washington, D.C: American Psychiatric Association, 2002.