

WHO antenatal care recommendations for a positive pregnancy experience"

Nutritional interventions update: zinc supplements during pregnancy



WHO antenatal care
recommendations for a positive
pregnancy experience"

**Nutritional interventions
update: zinc supplements
during pregnancy**



WHO antenatal care recommendations for a positive pregnancy experience. Nutritional interventions update: zinc supplements during pregnancy

ISBN 978-92-4-003046-6 (electronic version)

ISBN 978-92-4-003047-3 (print version)

© **World Health Organization 2021**

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: "This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition".

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

Suggested citation. WHO antenatal care recommendations for a positive pregnancy experience. Nutritional interventions update: zinc supplements during pregnancy. Geneva: World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IGO.

Cataloguing-in-Publication (CIP) data. CIP data are available at <http://apps.who.int/iris>.

Sales, rights and licensing. To purchase WHO publications, see <http://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <http://www.who.int/about/licensing>.

Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Editing, design and layout by Green Ink.

Contents

Acknowledgements	iv
Acronyms and abbreviations	v
Executive summary	vi
1 Introduction	1
2 Methods	3
3 Evidence and recommendation on antenatal zinc supplementation	7
4 Dissemination and implementation of the recommendation	15
5 Research implications	16
6 Updating the guideline	17
7 References	18
Annex 1. External experts and WHO staff involved in the preparation of the guideline	20
Annex 2. Summary of declarations of interest from the Guideline Development Group members, observers and External Review Group members and how they were managed	23
Annex 3. Zinc supplementation: GRADE tables	25

Acknowledgements

The Department of Sexual and Reproductive Health and Research (SRH), the Department of Nutrition and Food Safety (NFS) and the Department of Maternal, Newborn, Child and Adolescent Health and Ageing (MCA) of the World Health Organization (WHO) gratefully acknowledge the contributions that many individuals and organizations have made to the updating of this guideline recommendation.

María Barreix, Maurice Bucagu, Doris Chou, Olufemi Oladapo, Lisa Rogers and Özge Tunçalp were the members of the WHO Steering Group who managed the guideline development process. The members of the Guideline Development Group (GDG) included Niveen Abu-Rmeileh, Atf Ghérissi, Gill Gyte, James Neilson, Lisa Noguchi, Nafissa Osman, Erika Ota, Robert Pattinson, Kathleen Rasmussen, Rusidah Selamat, Harshpal Singh Sachdev and Charlotte Warren. James Neilson served as chair of the GDG. We thank members of the External Review Group (ERG), including Lorena Binfa, Tamar Kabakian, Bhavya Reddy and Petr Velebil.

We would also like to thank the authors of the updated Cochrane systematic review used, for their collaboration, and Frances Kellie and Leanne Jones, who facilitated this collaboration process. Soo Downe and Kenny Finlayson performed the qualitative reviews that informed the values, acceptability and feasibility criteria of the evidence-to-decision framework and graded the qualitative evidence for the WHO antenatal care guideline, which were also employed for this update. Therese Dowswell and Theresa Lawrie, with members of the WHO Steering Group, synthesized and reviewed the evidence and drafted the evidence-to-decision framework and the final guideline document.

We acknowledge the various organizations that were represented by observers at the technical consultation, including: Nita Dalmiya of the United Nations Children's Fund (UNICEF); Alison Tumilowicz of the Bill & Melinda Gates Foundation; Florence West of the International Confederation of Midwives (ICM); Petra ten Hoope-Bender of the United Nations Population Fund (UNFPA); Lindy Fenlason and Mary Ellen Stanton of the United States Agency for International Development (USAID). We appreciate the contributions of WHO Regional Office staff to this update: Nino Berdzuli, Karima Gholbzouri, Rodolfo Gomez, Anoma Jayathilaka, Léopold Ouedraogo and Howard Sobel.

Funding was provided for this updated recommendation by USAID and the United Nations Development Programme (UNDP)-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP). Donors do not fund specific guidelines and do not participate in any decision related to the guideline development process, including the composition of research questions, membership of the guideline groups, conduct and interpretation of systematic reviews, or formulation of recommendations.

Acronyms and abbreviations

ANC	antenatal care
CI	confidence interval
DECIDE	Developing and Evaluating Communication strategies to Support Informed Decisions and Practice based on Evidence
DOI	declaration of interest
ERG	External Review Group
EtD	evidence-to-decision
GDG	Guideline Development Group
GRADE	Grading of Recommendations Assessment, Development and Evaluation
GRADE-CERQual	Confidence in the Evidence from Reviews of Qualitative research
GSG	Guideline Steering Group
HRP	UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction
ICM	International Confederation of Midwives
LMICs	low- and middle-income countries
MCA	Department of Maternal, Newborn, Child and Adolescent Health and Ageing
NFS	Department of Nutrition and Food Safety
RCT	randomized controlled trial
RR	risk ratio
SRH	Department of Sexual and Reproductive Health and Research
UNDP	United Nations Development Programme
UNFPA	United Nations Population Fund
UNICEF	United Nations Children's Fund
US\$	United States dollar
USAID	United States Agency for International Development
WHO	World Health Organization

Executive summary

Introduction

Evidence from a systematic review on antenatal zinc supplementation was evaluated as part of the World Health Organization (WHO) antenatal care (ANC) guideline development process in 2016, and the following recommendation on zinc supplementation was made: “Zinc supplementation for pregnant women is only recommended in the context of rigorous research.” The Guideline Development Group (GDG) made this recommendation because it felt that the evidence on the intervention was incomplete and that more research was necessary.

Since the publication of the systematic review, four additional randomized controlled trials have been published; therefore, in April 2019, the Executive Guideline Steering Group (GSG) prioritized the WHO recommendation on antenatal zinc supplementation for updating.

Zinc is a trace element found in many foods, particularly in meat, but also in dairy products, legumes and unrefined cereals. It plays an important role in many biological processes that contribute to human growth and development, and also to immunity. As it is not stored in the body, it needs to be consumed regularly to prevent zinc deficiency, which is particularly common in low- and middle-income countries, where dietary intake of zinc-rich foods is often low. However, the effects of zinc deficiency in pregnancy have not been clearly established.

In January 2021, a WHO-convened GDG comprising most of the 2016 GDG members re-evaluated the evidence on antenatal zinc supplementation, updating the recommendation on zinc in accordance with WHO’s living guidelines approach.

Target audience

The target audience of this updated recommendation includes national and local public health policy-makers, implementers and managers of national and local maternal and child health programmes, concerned nongovernmental and other organizations, professional societies involved in the planning and management of maternal and child health services, health professionals (including obstetricians, midwives, nurses, nutritionists and general medical practitioners) and academic staff involved in training health professionals.

Guideline development methods

The updating of this recommendation was guided by the standardized operating procedures described in the *WHO handbook for guideline development*. These involve: (i) identification of the priority question and outcomes (done as part of the ANC guideline development process); (ii) evidence retrieval and synthesis; (iii) assessment of the evidence; (iv) formulation of the recommendation; and (v) planning for the dissemination, implementation, impact evaluation and updating of the recommendation. The scientific evidence supporting the recommendation was synthesized using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) and Confidence in the Evidence from Reviews of Qualitative research (GRADE-CERQual) approaches, for quantitative and qualitative evidence, respectively. An up-to-date systematic review was used to prepare an evidence profile for the recommendation prioritized for updating. The DECIDE (Developing and Evaluating Communication Strategies to Support Informed Decisions and Practice based on Evidence) framework, an evidence-to-decision tool that includes intervention effects, values, resources, equity, acceptability and feasibility criteria, was used to guide the formulation and approval of the recommendation by the GDG, an international group of experts that was convened for this process during an online GDG meeting on 13 January 2021. For consistency and continuity, the GDG, including the chair, comprised most of the same members as the ANC guideline GDG.

Recommendations

The WHO meeting led to the retention of the 2016 recommendation on antenatal zinc supplementation (Table 1). The GDG had the option to recommend the intervention, not recommend the intervention or recommend the intervention under certain conditions (in specific contexts, with targeted monitoring and evaluation, in the context of rigorous research). The GDG experts also provided additional remarks that they considered necessary

in the understanding and implementation of the recommendation. Users of the guideline should refer to these remarks, as well as to the evidence summary, for further information about the basis of this WHO recommendation.

This recommendation applies to pregnant women and adolescent girls within the context of routine ANC.

Table 1. The WHO recommendation on antenatal zinc supplementation for a positive pregnancy experience

Zinc supplementation for pregnant women is recommended only in the context of rigorous research.
Remarks
<ul style="list-style-type: none">• The Guideline Development Group agreed to retain the WHO recommendation found in the 2016 WHO antenatal care (ANC) guideline.• WHO does not recommend zinc supplementation as part of routine ANC. It is recommended in the context of rigorous research to improve our knowledge of its effect on maternal and newborn health outcomes. Research is particularly needed on how zinc status is impacted by other nutritional supplementation (e.g. iron and/or calcium) given as part of routine ANC. Additionally, research is needed on the efficacy of zinc supplementation, provided either alone or with other nutritional supplements (e.g. iron and folic acid, calcium, according to national guidelines/standard of care), on maternal and neonatal outcomes. Multiple doses of zinc, iron and/or calcium may need to be evaluated based on the current national standard of care. Research on the effectiveness or the implementation of zinc supplementation is not identified as a priority at this time.• Pregnant women should be encouraged and supported to receive adequate nutrition, which is best achieved through consumption of a healthy, balanced diet, and to refer to guidelines on healthy eating.

1 Introduction

1.1 Background

Zinc supplementation in pregnancy

Zinc is a trace element found in many foods, particularly in meat, but also in dairy products, legumes and unrefined cereals; diets low in bioavailable zinc are those in which animal protein is low and intake of cereals is high (1). Zinc is not stored in the body, so physiological needs must be met by dietary intake alone (2). In pregnant women, the average physiological requirement of zinc is estimated to double in the third trimester and almost triple during lactation (2). Therefore, many pregnant women are potentially at risk of zinc deficiency through inadequate dietary intake, exacerbated by the increased nutritional demands of pregnancy and lactation. In addition, routine iron supplementation may prevent women from meeting their zinc requirements by competing with zinc for absorption (3). This may also occur with calcium supplements and foods fortified with inorganic calcium salts (1).

Zinc plays an important role in many biological processes that contribute to human growth and development. Among children, the prevalence of stunting has been used as an indicator of zinc deficiency (1, 3). Zinc deficiency increases susceptibility to infections, and supplementing zinc among children in disadvantaged populations has been shown to reduce the incidence of diarrhoea and pneumonia (1–5). The World Health Organization (WHO) and the United Nations Children’s Fund (UNICEF) recommend zinc supplementation is given along with oral rehydration salts for the clinical management of diarrhoea in children (6).

In pregnancy, zinc deficiency has been linked to prolonged labour, postpartum haemorrhage, pre-eclampsia, preterm birth and post-term pregnancies (7); however, these associations have not been well established. This evidence-to-decision (EtD) framework presents research evidence on the effects and other considerations relevant to recommendations on antenatal zinc supplements compared with no zinc supplements or placebo.

The updated recommendation in the context of the WHO antenatal care guideline

In 2016, the following recommendation on antenatal zinc supplementation was made: “Zinc supplementation for pregnant women is only recommended in the context of rigorous research”(8). The Guideline Development Group (GDG) made this recommendation because it felt that the evidence on the intervention was incomplete and that more research was necessary. Since the publication of the 2016 WHO antenatal care (ANC) guideline, the systematic review, on which the 2016 recommendation was based, has been updated to include four additional trials (9); hence the need to re-evaluate the evidence.

1.2 Rationale and objectives

As part of the WHO’s normative work on supporting evidence-informed policies and practices and its living guidelines approach (10), the Department of Sexual and Reproductive Health and Research (SHR), the Department of Maternal, Newborn, Child and Adolescent Health and Ageing (MCA) and the Department of Nutrition and Food Safety (NFS) prioritized the updating of this recommendation on zinc supplements following the advice of the Executive Guideline Steering Group (GSG) in response to the identification of updated evidence on this intervention.

1.3 Target audience

The recommendation in this global guideline is intended to inform the development of relevant national and local health policies and clinical protocols. Therefore, the target audience of this guideline includes national and local public health policy-makers, implementers and managers of national and local maternal and child health programmes, concerned nongovernmental and other organizations, professional societies involved in the planning and management of maternal and child health services, health professionals (including obstetricians, midwives, nurses, nutritionists and general medical practitioners), researchers, and academic staff involved in training health professionals.

1.4 Scope of the recommendations

This updated recommendation is relevant to all pregnant women and adolescent girls receiving ANC in any health-care facility or community-based setting and to their fetuses and newborns. The guideline question was prioritized during the WHO 2016 ANC guideline development process. In 2019, the recommendation was prioritized for updating in the context of WHO's living guideline commitment (10). The outcomes of interests are, therefore, the same as those prioritized for the ANC guideline relevant to nutritional interventions (Box 1).

Box 1: Outcomes of interest in antenatal care nutritional interventions

Maternal outcomes	Fetal/neonatal outcomes
Infections	Neonatal infections
Anaemia	Small for gestational age
Pre-eclampsia/eclampsia	Low birthweight
Gestational diabetes mellitus	Preterm birth
Mode of delivery	Congenital anomalies
Excessive weight gain	Macrosomia (large for gestational age)
Side-effects	Fetal/neonatal mortality
Maternal mortality	
Maternal satisfaction	

2 Methods

This recommendation is an update of one of 49 recommendations that were published in *WHO recommendations on antenatal care for a positive pregnancy experience* (8). The recommendation was developed initially using the standardized operating procedures described in the *WHO handbook for guideline development* (11). In summary, the process included: (i) identification of priority question and outcomes, (ii) retrieval of evidence, (iii) assessment and synthesis of the evidence, (iv) formulation of recommendation, and (v) planning for the implementation, dissemination, impact evaluation and updating of the recommendation. This recommendation was identified by the Executive GSG as a high priority for updating in response to new evidence on this question.

2.1 Contributors to the guideline

Executive Guideline Steering Group

The Executive GSG is an independent panel of external experts and relevant stakeholders from the six WHO regions. This group advises WHO on the prioritization of new and existing questions in maternal and perinatal health for recommendation development or updating.

WHO Steering Group

The WHO Steering Group that managed the updating process comprised the same staff members from the Departments of SRH, MCA and NFS as the WHO ANC guideline of 2016 (see Annex 1 for the list of members). The WHO Steering Group drafted the priority question in PICO (population, intervention, comparator, outcome) format and identified individuals to be invited to participate as guideline methodologists, and in the guideline development and external review groups. In addition, the WHO Steering Group supervised the evidence retrieval and synthesis, organized the technical consultation and finalized the guideline document. Additionally, WHO regional office representatives for sexual and reproductive health were invited to all GDG sessions and provided comments on the updated recommendation document. The WHO Steering Group in collaboration with WHO regional offices will oversee the dissemination of the updated recommendation.

Guideline Development Group

The WHO Steering Group identified and invited 12 external experts and stakeholders from the six WHO regions to constitute the GDG, ensuring representation, gender balance and no important conflicts of interest. These experts also served in the GDG for the WHO ANC guideline's nutrition recommendations of 2016. They were a diverse group of individuals with expertise in research, guideline development methods and clinical policy and programmes relating to ANC interventions, plus a patient/consumer representative. The GDG appraised the evidence used to inform the recommendation, advised on the interpretation of this evidence and formulated the final recommendation during an online GDG meeting on 13 January 2021. In addition, GDG members reviewed and approved the final guideline document before its submission to the WHO Guidelines Review Committee for approval. A list of the GDG members can be found in Annex 1.

External Review Group

The External Review Group (ERG) was a geographically and gender-balanced group with no important conflicts of interest (see Annex 2). There were four members, including technical experts and other stakeholders with interests in the provision of evidence-informed ANC. This group peer-reviewed the draft version of the guideline document to identify any factual errors and comment on the clarity of the language, contextual issues and implications for implementation. The group ensured that the guideline decision-making processes had considered and incorporated the contextual values and preferences of people affected by the recommendation, including pregnant women and adolescent girls, health-care professionals and policy-makers. It was not within the ERG's remit to change the recommendation formulated by the GDG.

Systematic review team and guideline methodologists

The managing editors of the Cochrane pregnancy and childbirth group coordinated the updating of the quantitative systematic review and facilitated collaboration between systematic review authors and guideline methodologists. Working closely with the WHO Steering Group, methodologists from The Evidence-based Medicine Consultancy in the United Kingdom of Great Britain and Northern Ireland appraised the quantitative evidence using WHO's standardized operating procedures for GRADE (Grading of Recommendations Assessment, Development and Evaluation) methodology (12). Two qualitative-evidence experts from the University of Central Lancashire in the United Kingdom systematically reviewed qualitative studies related to women's and health professionals' views on ANC, and synthesized this evidence.

External partners and observers

Six representatives of the ICM, UNFPA, USAID, UNICEF and the Bill & Melinda Gates Foundation were invited to the final GDG meeting to serve as observers. All these organizations are potential implementers of the proposed guideline with a history of collaboration with the WHO in guideline dissemination and implementation. Observers do not participate in the formulation of recommendations.

2.2 Declaration of interests by external contributors

WHO requires that experts serving in an advisory role disclose any circumstances that could give rise to actual or ostensible conflicts of interest. In accordance with the *WHO guidelines for declaration of interests (WHO Experts)* (13), all GDG members and other external collaborators were asked to declare in writing any competing interests (whether academic, financial or other) at the time of the invitation to participate in the ANC guideline development process. The standard WHO form for declaration of interest (DOI) was completed and signed by each expert. The WHO Steering Group reviewed all the DOI forms before finalizing experts' invitations to participate. Where any conflicts of interest were declared, the WHO Steering Group determined whether they were serious enough to affect the individual's ability to make objective judgements about the evidence or recommendation. To ensure consistency, the WHO Steering Group applied the criteria for assessing the severity of a conflict of interest in the *WHO handbook for guideline development* (11).

All findings from DOI statements were managed in accordance with the WHO DOI guidelines on a case-by-case basis and communicated to the experts. Where a conflict of interest was not considered significant enough to pose any risk to the guideline development process or reduce its credibility, the expert was only required to declare such a conflict at the GDG meeting, and no further action was taken. A summary of the DOI statements and information on how conflicts of interest were managed are included in Annex 2. To strengthen public trust and transparency in connection with WHO meetings involving the provision of expert advice in developing technical norms and standards, the names and brief biographies of individuals considered for participation in this guideline together, with a description of the objectives of relevant meetings, were made public ahead of the planned online GDG meeting, to allow time for public notice and comment.

2.3 Identifying priority questions and outcomes

The priority question and outcomes were aligned with those of the ANC guideline. This question and outcomes were originally informed through an extensive scoping exercise of existing clinical practice guidelines relevant to routine ANC, supplemented by searching the Cochrane Database of Systematic Reviews for existing key systematic reviews relevant to ANC. Critical and important outcomes were informed by these reviews, and by a WHO-commissioned scoping qualitative review of what women want during pregnancy (14). The findings of the latter revealed that pregnant women want a positive pregnancy experience, defined as maintaining physical and sociocultural normality; maintaining a healthy pregnancy and baby; having an effective transition to positive labour and birth; and achieving a positive motherhood. This composite outcome of a positive pregnancy experience became the overarching principle of ANC guideline recommendations.

2.4 Evidence identification and retrieval

Evidence to support this recommendation was derived from a number of sources by the methodologists working closely with the WHO Steering Group. An updated systematic review was the primary source of evidence on the effectiveness of oral antenatal zinc supplementation. Earlier versions of this review – in which evidence on effectiveness was derived from randomized controlled trial (RCT) data assessed and synthesized using standardized Cochrane methodology – supported the ANC guideline recommendation of 2016. The up-to-date RevMan file was retrieved from the Cochrane pregnancy and childbirth group and customized to reflect the key comparisons, GDG-specified subgroup analyses, and outcomes relevant to the ANC guideline. Evidence was evaluated according to standard operating procedures approved by the WHO Steering Group, and evidence profiles (in the form of GRADE tables) were prepared, including assessments of the certainty of the evidence, for the comparisons of interest.

Two qualitative systematic reviews commissioned by the WHO Steering Group for the 2016 guideline development process informed the values, acceptability and feasibility criteria of the EtD frameworks to inform the recommendation (see below for more information, under section 2.6 – Preparation of the evidence summary) (14, 15). Additionally, systematic reviews of cost-effectiveness were sought through PubMed searches of the literature.

2.5 Quality assessment and grading of the evidence

The GRADE approach (12) was used to appraise the certainty of quantitative evidence, meaning that the certainty of evidence for each outcome was rated as “high”, “moderate”, “low”, or “very low” based on a set of established criteria. As a baseline, the evidence from the systematic reviews was rated “high certainty” because it was derived from RCTs; this rating was then downgraded according to considerations of risk of bias, inconsistency, imprecision, indirectness and publication bias or other considerations.

Qualitative evidence was derived from a qualitative evidence synthesis performed for the WHO 2016 ANC guideline (14, 15). Previously subjected to a quality appraisal using the GRADE-CERQual (Confidence in the Evidence from Reviews of Qualitative research) tool, the evidence was not regraded for this updated recommendation. The GRADE-CERQual tool, which uses a similar approach conceptually to other GRADE tools, rates the level of confidence that can be placed in qualitative evidence synthesis according to four components: methodological limitations of the individual studies, adequacy of data and coherence and relevance of the findings to the review question (16).

2.6 Preparation of the evidence summary

The WHO Steering Group supervised and finalized the preparation of the evidence summary and profile, in collaboration with the guideline methodologists, using the DECIDE (Developing and Evaluating Communication strategies to Support Informed Decisions and Practice based on Evidence) framework. DECIDE is an EtD tool that includes explicit and systematic consideration of research evidence on interventions according to six criteria, namely effects, values, resources, equity, acceptability and feasibility (17). These six EtD criteria were populated with the research evidence, where available; in addition, information from other sources was described in the additional considerations subsections of each criterion. The certainty of the graded evidence on intervention effectiveness was systematically interpreted in EtD frameworks according to guidance from the Cochrane group on effective practice and organization of care (18).

2.7 Formulation of the recommendation

GDG members and other participants were provided with the evidence summary in advance of the online GDG meeting held on 13 January 2021, organized by the WHO Steering Group from Geneva, Switzerland. During the technical consultation, under the leadership of the GDG chair, the GDG members reviewed, discussed and made judgements on the impact of the intervention for each of the EtD criteria. GDG judgements were summarized in a table, before finalizing the recommendation and remarks. The intervention could be recommended, not recommended, or recommended only under certain conditions (in specific contexts, with targeted monitoring and evaluation, in the context of rigorous research).

2.8 Decision-making process

The online GDG meeting was guided by a clear protocol, designed to allow the recommendation to be formulated through a process of group discussion, until consensus was reached. The final adoption of the recommendation and, if necessary, the context in which the recommendation would apply were confirmed by unanimous consensus (i.e. full agreement among all GDG members).

2.9 Guideline preparation and peer review

Following the online GDG meeting, members of the WHO Steering Group, assisted by methodologists, drafted a full guideline document to accurately reflect the deliberations and decisions of participants. A preliminary version of the document was sent electronically to the participants and the ERG for final review and technical comments. The WHO Steering Group carefully evaluated the input of the peer reviewers for inclusion in the guideline document and made revisions to the guideline draft as needed. After the GDG meetings and peer review process, further modifications to the guideline by the WHO Steering Group were limited to corrections of factual errors and improvements in language to address any lack of clarity. The document was then submitted for executive clearance according to established WHO publication procedures.

3 Evidence and recommendation on antenatal zinc supplementation

This section provides the WHO recommendation adopted by the GDG on antenatal zinc supplementation with its corresponding evidence summary. Evidence on the effectiveness of zinc supplementation is further detailed in GRADE tables in Annex 3. To ensure that the recommendation is correctly understood, additional remarks reflecting the summary of the discussion by the GDG are included below the recommendation.

WHO recommendation on antenatal zinc supplementation

Zinc supplementation for pregnant women is recommended only in the context of rigorous research.

Remarks

- This Guideline Development Group agreed to retain the WHO recommendation found in the 2016 WHO antenatal care guideline (8).
- WHO does not recommend zinc supplementation as part of routine antenatal care. It is recommended only in the context of rigorous research to improve our knowledge of its effect on maternal and newborn health outcomes. Research is particularly needed on how zinc status is impacted by other nutritional supplementation (e.g. iron and/or calcium) given as part of routine antenatal care. Additionally, research is needed on the efficacy of zinc supplementation – provided either alone or with other nutritional supplements (e.g. iron, folic acid and calcium, according to national guidelines/standard of care) – on maternal and neonatal outcomes. Multiple doses of zinc, iron and/or calcium may need to be evaluated based on the current national standard of care. Research on the effectiveness or the implementation of zinc supplementation is not identified as a priority at this time.
- Pregnant women should be encouraged and supported to receive adequate nutrition, which is best achieved through a healthy, balanced diet, and to refer to guidelines on healthy eating (19).

3.1 The priority question

For pregnant women (population), does zinc supplementation (intervention) compared with no zinc supplementation (comparator) improve maternal and perinatal health outcomes (outcome)?

3.2 Assessment

Effects of the intervention

What are the anticipated effects of antenatal zinc supplements compared with no zinc supplements (or placebo)?

Research evidence

The following evidence was derived from a systematic review (9) that updated evidence from a review published in 2015 (7). The updated review includes data from four additional trials, so that there was a total of 25 RCTs, involving more than 18 000 pregnant women. Studies were carried out in Bangladesh (two studies), Chile, China, Denmark, Egypt, Ghana, Indonesia (two studies), the Islamic Republic of Iran (three studies), Nepal, Pakistan, Peru (two studies), South Africa, United Republic of Tanzania, the United Kingdom (three studies) and the United States of America (four studies). The dose of daily zinc supplementation ranged from 5 mg to 50 mg of zinc per day, although in one trial some women received up to 90 mg of zinc per day. Eleven trials compared zinc supplementation with placebo. There was a wide variation in the size of trials (56 women recruited in the smallest, 4926 women in the largest) and in women's nutritional and zinc status at trial entry. Women's gestational age at recruitment and the duration of supplementation also varied across trials (before conception in one trial, in the first or second trimester in most trials, after 26 weeks' gestation in two trials, and up to 6 months postpartum in one trial), as did compliance with treatment.

Maternal outcomes

Pre-eclampsia: The evidence suggests that zinc supplementation may make little or no difference to pre-eclampsia compared with no zinc supplements (six trials, 2568 women; risk ratio, RR, 0.93, 95% confidence interval, CI, 0.62 to 1.42; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Maternal infection: The evidence suggests that zinc supplementation may make little or no difference to maternal infection compared with no zinc supplements (four trials, 1891 women; RR 0.94, 95% CI 0.72 to 1.23; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Side-effects: Side-effects were not generally reported in trials.

The evidence on caesarean section was of very low certainty and there were no relevant data on maternal anaemia, maternal mortality, gestational diabetes mellitus or positive pregnancy experience.

Fetal/neonatal outcomes

Small for gestational age: Zinc supplementation probably makes little or no difference to the small for gestational age outcome compared with no zinc supplements (nine trials, 5330 babies; RR, 1.02, 95% confidence interval, CI, 0.92 to 1.12; *moderate-certainty evidence, downgraded due to design limitations*).

Low birthweight: Zinc supplementation probably makes little or no difference to low birthweight compared with no zinc supplements (17 trials, 7399 babies; RR 0.94, 95% CI 0.79 to 1.13; *moderate-certainty evidence, downgraded due to design limitations*).

Preterm birth: Zinc supplementation probably makes little or no difference to preterm birth compared with no zinc supplements (20 trials, 9454 babies; RR 0.90, 95% CI 0.78 to 1.04; *moderate-certainty evidence, downgraded due to design limitations*).

Neonatal mortality: Zinc supplementation may make little or no difference to neonatal mortality compared with no zinc supplements (three trials, 1965 babies; RR 2.24, 95% CI 0.40 to 14.83; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Stillbirth: Zinc supplementation may make little or no difference to stillbirth compared with no zinc supplements (six trials, 2898 babies; RR 1.32, 95% CI 0.85 to 2.04; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Perinatal mortality: Zinc supplementation may make little or no difference to perinatal mortality compared with no zinc supplements (two trials, 2489 babies; RR 1.10, 95% CI 0.81 to 1.51; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Congenital anomalies: Zinc supplementation may make little or no difference to congenital malformation compared with no zinc supplements (five trials, 1106 babies; RR 0.67, 95% CI 0.33 to 1.35; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Neonatal infection: Zinc supplementation may make little or no difference to neonatal sepsis compared with no zinc supplements (two trials, 736 babies; RR 0.17, 95% CI 0.03 to 1.01; *low-certainty evidence, downgraded due to design limitations and imprecision*).

Summary of effects

Low-certainty evidence suggests that zinc supplementation may have little or no effect on pre-eclampsia and maternal infections; there is insufficient evidence available for other maternal outcomes. In addition, low-certainty evidence suggests that supplementation may have little or no effect on any of the important fetal and neonatal outcomes.

Additional considerations

- WHO recommends iron and folic acid supplementation containing 30 mg to 60 mg of iron and 0.4 mg of folic acid for all pregnant women, and calcium supplementation for women with low dietary intake and those at risk of pre-eclampsia; these supplements may potentially reduce the bioavailability of zinc (1).

Desirable effects

How substantial are the desirable anticipated effects of zinc supplements compared with no zinc supplements?

Judgement					
<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Trivial	<input type="checkbox"/> Small	<input type="checkbox"/> Moderate	<input type="checkbox"/> Large

Rationale for judgement: There appears to be no clear improvement in pregnancy outcomes.

Undesirable effects

How substantial are the undesirable anticipated effects of zinc supplements compared with no zinc supplements?

Judgement					
<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Trivial	<input type="checkbox"/> Small	<input type="checkbox"/> Moderate	<input type="checkbox"/> Large

Rationale for judgement: There was no evidence suggesting harm but adverse events were generally not reported in the studies.

Certainty of the evidence

What is the overall certainty of the evidence of effects of zinc supplements compared with no zinc supplements?

Judgement				
<input type="checkbox"/> No included studies	<input type="checkbox"/> Very low	<input checked="" type="checkbox"/> Low	<input type="checkbox"/> Moderate	<input type="checkbox"/> High

Rationale for judgement: Low certainty was the most common GRADE rating.

Values

Is there important uncertainty about, or variability in, how much women (and their families) value the main outcomes associated with zinc supplements?

A scoping review of what women want from ANC informed the outcomes for the WHO ANC guideline (14). Evidence showed that women from various resource settings valued having a positive pregnancy experience comprising three equally important components, namely effective clinical practices (interventions and tests), relevant and timely information, and psychosocial and emotional support, each provided by practitioners with good clinical and interpersonal skills within a well-functioning health system (*high confidence in the evidence*).

Judgement			
<input type="checkbox"/> Important uncertainty or variability	<input type="checkbox"/> Possibly important uncertainty or variability	<input checked="" type="checkbox"/> Probably no important uncertainty or variability	<input type="checkbox"/> No important uncertainty or variability

Rationale for judgement: It is important to pregnant women that clinical practices are effective.

Balance of effects

Does the balance between desirable and undesirable effects favour zinc supplements or no zinc supplements?

Judgement						
<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Favours no zinc supplements	<input type="checkbox"/> Probably favours no zinc supplements	<input checked="" type="checkbox"/> Does not favour zinc supplements or no zinc supplements	<input type="checkbox"/> Probably favours zinc supplements	<input type="checkbox"/> Favours zinc supplements

Rationale for judgement: The low-certainty evidence suggests that there are neither benefits nor harms with zinc supplements.

3.3 Resources

How large are the resource requirements (costs)?

Research evidence

No research evidence on the costs and cost-effectiveness of implementing zinc supplementation was found.

Main resource requirements

The main resource requirement is probably the cost of the supplements. The UNICEF Supply Catalogue pricing accessed on 3 June 2021 gave an indicative price of US\$ 1.34 for a blister pack of 100 tablets of 20 mg each, which is equivalent to US\$ 3.20 for six-month supply (20). However, to ensure that women received relevant and accurate information about the supplements, staff training would be required, which would also have cost implications.

Additional considerations

- The cost-effectiveness of zinc supplements for pregnant women is uncertain because the systematic review found no strong evidence of their effectiveness in relation to the pregnancy outcomes evaluated.
- To have an impact on child mortality, it has been suggested that food fortification may be the most cost-effective strategy to supplement zinc among populations with a high prevalence of zinc deficiency (3).

Resources required

How costly are the resources required for zinc supplements compared with no zinc supplements?

Judgement						
<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Large costs	<input checked="" type="checkbox"/> Moderate costs	<input type="checkbox"/> Negligible costs or savings	<input type="checkbox"/> Moderate savings	<input type="checkbox"/> Large savings

Rationale for judgement: The cost of zinc supplements is similar to that of iron and folic acid supplements, and, if implemented, may double the cost of micronutrient supplements provided to women during pregnancy.

Certainty of evidence on required resources

What is the certainty of the evidence on costs?

Judgement				
<input type="checkbox"/> No included studies	<input type="checkbox"/> Very low	<input type="checkbox"/> Low	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> High

Rationale for judgement: The supply costs are taken from the UNICEF Supply Catalogue.

Cost-effectiveness

How cost-effective are zinc supplements compared with no zinc supplements?

Judgement						
<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Favours no zinc supplements	<input type="checkbox"/> Probably favours no zinc supplements	<input type="checkbox"/> Does not favour zinc supplements or no zinc supplements	<input type="checkbox"/> Probably favours zinc supplements	<input type="checkbox"/> Favours zinc supplements

Rationale for judgement: The evidence did not show zinc supplementation to be effective, so cost-effectiveness cannot be evaluated.

3.4 Equity

What would be the impact of zinc supplements compared with no zinc supplements on health equity?

Research evidence

None.

Additional considerations
<ul style="list-style-type: none"> Nutritional deficiencies are common in low- and middle-income countries (LMICs). Effective interventions to improve the nutritional status of pregnant women and girls in these settings could help to address health inequalities by preventing illnesses related to vitamin and mineral deficiencies.

Judgement						
<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Reduced	<input type="checkbox"/> Probably reduced	<input type="checkbox"/> Probably no impact	<input checked="" type="checkbox"/> Probably increased	<input type="checkbox"/> Increased

Rationale for judgement: There is no clear evidence of the effectiveness of zinc supplementation for pregnancy outcomes; the effectiveness of zinc supplementation on general maternal health in LMICs was not a prioritized guideline outcome.

3.5 Acceptability

Would zinc supplements be acceptable to key stakeholders?

Research evidence

A systematic review of qualitative research exploring women’s views and experiences of ANC suggests that women tend to view ANC as a source of knowledge and information and generally appreciate any advice (including dietary or nutritional) that may lead to a healthy baby and a positive pregnancy experience (*high confidence in the evidence*) (15).

The same review explored health professionals’ views of ANC, and suggests that health professionals are keen to offer general health-care advice and specific pregnancy-related information (*low confidence in the evidence*) but sometimes feel they do not have the appropriate training, and the resources and time to deliver the service in the informative, supportive and caring manner that women want (*high confidence in the evidence*) (15).

Additional considerations

None.

Judgement

<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> No	<input checked="" type="checkbox"/> Probably No	<input type="checkbox"/> Probably Yes	<input type="checkbox"/> Yes
-------------------------------------	---------------------------------	-----------------------------	---	---------------------------------------	------------------------------

Rationale for judgement: If zinc supplementation improved pregnancy outcomes, supplementation would probably be acceptable. However, as there is no clear evidence of effectiveness, it may not be acceptable.

3.6 Feasibility

Would zinc supplementation be feasible to implement?

Research evidence

Evidence derived from a qualitative evidence synthesis conducted to support the WHO ANC guideline development shows that where there are likely to be additional costs associated with supplementation (*high confidence in the evidence*), women may be less likely to engage with services (15). In addition, in a number of LMIC settings, providers felt that a lack of resources, both in terms of the availability of recommended supplements and the lack of suitably trained staff to deliver nutritional information, was an issue, which may limit the implementation of this intervention (*high confidence in the evidence*).

Additional considerations

- On the demand side, if zinc supplements are free and available, routine supplementation may be feasible. On the supply side, however, there may be several considerations to take into account, such as changes in regulatory norms and policies (e.g. tariffs, labelling, imports, government oversight, etc.), how sustainable the production is (local or imported) and how to guarantee product availability.
- In addition, the lack of effectiveness of this intervention suggests that it would not be feasible in LMICs where health-care expenditure is constrained.

Judgement

<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> No	<input checked="" type="checkbox"/> Probably No	<input type="checkbox"/> Probably Yes	<input type="checkbox"/> Yes
-------------------------------------	---------------------------------	-----------------------------	---	---------------------------------------	------------------------------

Rationale for judgement: There is no clear evidence of effectiveness.

3.7 Summary of GDG judgements on antenatal zinc supplementation (18)

Desirable effects	<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies		<input type="checkbox"/> Trivial	<input type="checkbox"/> Small	<input type="checkbox"/> Moderate	<input type="checkbox"/> Large
Undesirable effects	<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies		<input type="checkbox"/> Large	<input type="checkbox"/> Moderate	<input type="checkbox"/> Small	<input type="checkbox"/> Trivial
Certainty of the evidence on effects	<input type="checkbox"/> No included studies			<input type="checkbox"/> Very low	<input checked="" type="checkbox"/> Low	<input type="checkbox"/> Moderate	<input type="checkbox"/> High
Values				<input type="checkbox"/> Important uncertainty or variability	<input type="checkbox"/> Possibly important uncertainty or variability	<input checked="" type="checkbox"/> Probably no important uncertainty or variability	<input type="checkbox"/> No important uncertainty or variability
Balance of effects	<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Favours no zinc supplements	<input type="checkbox"/> Probably favours no zinc supplements	<input checked="" type="checkbox"/> Does not favour zinc supplements or no zinc supplements	<input type="checkbox"/> Probably favours zinc supplements	<input type="checkbox"/> Favours zinc supplements
Resources required	<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Large costs	<input checked="" type="checkbox"/> Moderate costs	<input type="checkbox"/> Negligible costs or savings	<input type="checkbox"/> Moderate savings	<input type="checkbox"/> Large savings
Certainty of evidence on required resources	<input type="checkbox"/> No included studies			<input type="checkbox"/> Very low	<input type="checkbox"/> Low	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> High
Cost-effectiveness	<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Favours no zinc supplements	<input type="checkbox"/> Probably favours no zinc supplements	<input type="checkbox"/> Does not favour zinc supplements or no zinc supplements	<input type="checkbox"/> Probably favours zinc supplements	<input type="checkbox"/> Favours zinc supplements
Equity	<input checked="" type="checkbox"/> Don't know	<input type="checkbox"/> Varies	<input type="checkbox"/> Reduced	<input type="checkbox"/> Probably reduced	<input type="checkbox"/> Probably no impact	<input type="checkbox"/> Probably increased	<input type="checkbox"/> Increased
Acceptability	<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies		<input type="checkbox"/> No	<input checked="" type="checkbox"/> Probably No	<input type="checkbox"/> Probably Yes	<input type="checkbox"/> Yes
Feasibility	<input type="checkbox"/> Don't know	<input type="checkbox"/> Varies		<input type="checkbox"/> No	<input checked="" type="checkbox"/> Probably No	<input type="checkbox"/> Probably Yes	<input type="checkbox"/> Yes

3.8 Conclusions

Recommendation

Zinc supplementation for pregnant women is recommended only in the context of rigorous research.

Judgement		
We do not recommend the intervention <input type="checkbox"/>	We recommend considering the intervention only <input type="checkbox"/> in specific contexts <input type="checkbox"/> with targeted monitoring and evaluation <input checked="" type="checkbox"/> in the context of rigorous research	We recommend the intervention <input type="checkbox"/>

Remarks

- This GDG agreed to retain the WHO recommendation found in the 2016 WHO ANC guideline (8).
- WHO does not recommend zinc supplementation as part of routine ANC. Zinc supplementation is recommended only in the context of rigorous research to improve our knowledge of its effect in pregnant women. Research is particularly needed on how zinc status is impacted by other nutritional supplementation (e.g. iron and/or calcium) given as part of routine ANC. Additionally, research is needed on the efficacy of zinc supplementation – provided either alone or with other nutritional supplements (e.g. iron and folic acid, calcium, according to national guidelines/standard of care) – on maternal and neonatal outcomes. Multiple doses of zinc, iron and/or calcium may need to be evaluated based on the current national standard of care. Research on the effectiveness or the implementation of zinc supplementation is not identified as a priority at this time.
- Pregnant women should be encouraged and supported to receive adequate nutrition, which is best achieved through a healthy, balanced diet, and to refer to guidelines on healthy eating (19).

Draft implementation considerations

Not applicable.

Research gaps

- The GDG agreed that more research on the efficacy of zinc supplementation in pregnancy was needed and that trials of zinc supplementation should consider other nutritional supplements (e.g. iron and/or calcium) that women may be receiving as part of routine ANC. As iron supplementation is recommended in all settings, and calcium supplementation is recommended in settings with low dietary intake as part of routine ANC, research is needed to evaluate the interactions among these nutrients in relation to maternal and newborn outcomes.
- In settings providing iron and folic acid as part of routine ANC, trials on zinc supplementation should include several doses of zinc along with either 30 mg and/or 60 mg of elemental iron, according to national guidance.
- Information should be recorded carefully on the timing of the different supplements, whether taken at the same time or different times of the day, along with an assessment of the feasibility and acceptability of taking multiple supplements at different times in these trials.

4 Dissemination and implementation of the recommendation

4.1 Recommendation dissemination

This updated global guideline will be available online for download and also as a printed publication. Online versions will be available on the WHO websites and other online platforms developed by the WHO Departments of SRH, NFS and MCA, and through the WHO ANC portal¹ and the WHO Reproductive Health Library and WHO e-Library of Evidence for Nutrition Actions.² Print versions will be distributed to WHO regional and country offices, ministries of health, WHO collaborating centres, nongovernmental organization partners, among others, using the same distribution list that was developed for the WHO ANC guideline. The updated recommendation and updated derivative products, in particular the *WHO Antenatal Care Recommendations Adaptation Toolkit* and its instruction manual, will be disseminated during meetings and scientific conferences attended by WHO staff (21). Social media channels will also be used. The executive summary and recommendation from this publication will be translated into the six United Nations languages for dissemination through the WHO regional offices and during meetings organized or attended by WHO staff.

4.2 Implementation considerations and applicability issues

This updated recommendation supersedes the WHO ANC guideline recommendation on zinc supplementation that was issued in 2016 (recommendation A6) (8). The GDG agreed that there were no new implementation considerations or applicability issues specific to this recommendation, as it was recommended in a research context. Monitoring of the uptake and impact of this updated recommendation will be integrated into that of the 2016 WHO ANC guideline (8). For implementation considerations related to *WHO recommendations on antenatal care for a positive pregnancy experience* in general, please refer to this guideline and associated derivative products, which are available on the WHO websites.

5 Research implications

The GDG agreed that more research on the efficacy of zinc supplementation in pregnancy is needed and that trials of zinc supplementation should consider other nutritional supplements (e.g. iron and/or calcium) that women may be receiving as part of routine ANC. As iron supplementation is recommended in all settings and calcium supplementation is recommended in settings with low dietary intake as part of routine ANC, research is needed to evaluate the interactions among these nutrients in relation to maternal and newborn outcomes.

In settings providing iron and folic acid as part of routine ANC, trials on zinc supplementation should include several doses of zinc along with 30 mg and/or 60 mg of elemental iron, according to national guidance.

Information should be recorded carefully on the timing of the different supplements, whether taken at the same time or different times of the day, along with an assessment of the feasibility and acceptability of taking multiple supplements at different times in these trials.

6 Updating the guideline

WHO convenes the Executive GSG biannually to review WHO's current portfolio of maternal and perinatal health recommendations, and to advise on the prioritization of new and existing questions for recommendation development and updating. WHO will monitor the publication of new randomized trials on this topic. Any concern about the validity of the recommendation will be promptly communicated via the guideline website,³ and plans will be made to update the recommendation, as necessary. WHO will prioritize its independent normative guidance informed by the strategic shifts embedded in its Constitution and the Thirteenth General Programme of Work 2019–2023.

All technical products developed during the process of developing this recommendation, including the Cochrane RevMan⁴ file customized for priority outcomes, and the basis for quality ratings within the GRADE process, will be archived in the departmental shared folder for future reference and use.

7 References

1. Vitamin and mineral requirements in human nutrition, second edition. Geneva: World Health Organization; 2004 (<https://www.who.int/nutrition/publications/micronutrients/9241546123/en/>, accessed 28 April 2021).
2. Trace elements in human nutrition and health. Geneva: World Health Organization; 1996 (<https://www.who.int/nutrition/publications/micronutrients/9241561734/en/>, accessed 28 April 2021).
3. Roohani N, Hurrell R, Kelishadi R, Schulin R. Zinc and its importance for human health: An integrative review. *J Res Med Sci.* 2013; 18(2):144-57.
4. Lassi ZS, Moin A, Bhutta ZA. Zinc supplementation for the prevention of pneumonia in children aged 2 months to 59 months. *Cochrane Database Syst Rev.* 2016; 12(12):CD005978. doi:10.1002/14651858.CD005978.pub3.
5. World Health Organization, United Nations University, United Nations Children's Fund. Composition of a multi-micronutrient supplement to be used in pilot programmes among pregnant women in developing countries. Geneva: WHO; 1999 (<https://apps.who.int/iris/handle/10665/75358>, accessed 28 April 2021).
6. World Health Organization, United Nations Children's Fund. Clinical management of acute diarrhoea. Geneva: WHO; 2004 (https://www.who.int/maternal_child_adolescent/documents/who_fch_cah_04_7/en/, accessed 28 April 2021).
7. Ota E, Mori R, Middleton P, Tobe-Gai R, Mahomed K, Miyazaki C, et al. Zinc supplementation for improving pregnancy and infant outcome. *Cochrane Database of Systematic Reviews.* 2015; 2:CD000230. doi:10.1002/14651858.CD000230.pub5.
8. WHO recommendations on antenatal care for a positive pregnancy experience. Geneva: World Health Organization; 2016 (<https://www.who.int/publications/i/item/9789241549912>, accessed 28 April 2021).
9. Carducci B, Keats EC, Bhutta ZA. Zinc supplementation for improving pregnancy and infant outcome. *Cochrane Database Syst Rev.* 2021; 3:CD000230. doi:10.1002/14651858.CD000230.pub6.
10. Vogel JP, Dowswell T, Lewin S, Bonet M, Hampson L, Kellie F, et al. Developing and applying a 'living guidelines' approach to WHO recommendations on maternal and perinatal health. *BMJ Global Health.* 2019; 4(4):e001683. doi:10.1136/bmjgh-2019-001683.
11. WHO handbook for guideline development, second edition. Geneva: World Health Organization; 2014 (<https://www.who.int/groups/guidelines-review-committee>, accessed 23 March 2021).
12. GRADE. Welcome to the GRADE working group [website]. The GRADE Working Group; 2021 (<https://www.gradeworkinggroup.org>, accessed 28 April 2021).
13. Declarations of interest. In: About WHO - Ethics [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/about/ethics/declarations-of-interest>, accessed 8 December 2020).
14. Downe S, Finlayson K, Tunçalp Ö, Gülmezoglu AM. What matters to women: a scoping review to identify the processes and outcomes of antenatal care provision that are important to healthy pregnant women. *BJOG.* 2016; 123(4):529-39. doi:10.1111/1471-0528.13819.
15. Downe S, Finlayson K, Tunçalp Ö, Gülmezoglu AM. Provision and uptake of routine antenatal services: a qualitative evidence synthesis. *The Cochrane database of systematic reviews.* 2019; 6(6):CD012392. doi:10.1002/14651858.CD012392.pub2.
16. Confidence in the Evidence from Reviews of Qualitative research [website]. GRADE-CERQual; 2018 (<https://cerqual.org/>, accessed 28 November 2020).
17. DECIDE GRADE. Key DECIDE tools [website]. 2015 (<https://www.decide-collaboration.eu>, accessed 28 April 2021).

18. Cochrane Effective Practice and Organisation of Care Group. Reporting the effects of an intervention in EPOC reviews. Oslo: Norwegian Knowledge Centre for the Health Services; 2018 (https://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/Resources-for-authors2017/how_to_report_the_effects_of_an_intervention.pdf, accessed 8 December 2020).
19. Healthy diet. Fact sheet No. 394. Geneva: World Health Organization; 2018 (https://www.who.int/nutrition/publications/nutrientrequirements/healthydiet_factsheet/en, accessed 16 November 2020).
20. UNICEF Supply Catalogue [website]. New York: United Nations Children's Fund; 2018 (<https://supply.unicef.org/>, accessed 22 October 2020).
21. Barreix M, Lawrie TA, Kidula N, Tall F, Bucagu M, Chahar R, et al. Development of the WHO antenatal care recommendations adaptation toolkit: a standardised approach for countries. *Health Res Policy Sys.* 2020;18:70. doi:10.1186/s12961-020-00554-4.

Annex 1. External experts and WHO staff involved in the preparation of the guideline

WHO Steering Group

María Barreix

Technical Officer
Department of Sexual and Reproductive Health and Research (SRH)
Maternal and Perinatal Health

Maurice Bucagu

Medical Officer
Department of Maternal, Newborn, Child and Adolescent and Health and Ageing
Policy, Planning and Programme Unit

Doris Chou

Medical Officer
Department of SRH
Maternal and Perinatal Health

Olufemi T. Oladapo

Unit Head
Department of SRH
Maternal and Perinatal Health

Lisa Rogers

Technical Officer
Department of Nutrition and Food Safety
Food and Nutrition Actions in Health Systems

Özge Tunçalp

Scientist
Department of SRH
Maternal and Perinatal Health

WHO regional office representatives

Regional Office for Europe

Nino Berdzuli

Regional Advisor
Director
Division of Country Health Programmes
Copenhagen, Denmark

Regional Office for Americas

Rodolfo Gomez

Reproductive Health Advisor
Latin American Center for Perinatology, Women and Reproductive Health
Montevideo, Uruguay

Regional Office for South East Asia

Anoma Jayathilaka

Medical Officer
Family Health, Gender and Life Course
New Delhi, India

Regional Office for the Eastern Mediterranean

Karima Gholbzouri

Regional Adviser
Reproductive & Maternal Health
Cairo, Egypt

Regional Office for Africa

Léopold Ouedraogo

Regional Adviser – focal point on sexual and reproductive health and rights
Reproductive, Maternal Health and Ageing
Brazzaville, Republic of the Congo

Regional Office of the Western Pacific

Howard Sobel

Regional Adviser
Maternal Child Health and Quality Safety
Manila, Philippines

Guideline Development Group

Niveen Abu-Rmeileh

Professor
Institute of Community and Public Health
Birzeit University
West Bank and Gaza Strip

Atf Ghérissi

International Expert
Health Sciences Education
Tunis, Tunisia

Gill Gyte

Consumer Editor
Cochrane Pregnancy and Childbirth Group
Liverpool, United Kingdom of Great Britain and Northern Ireland (United Kingdom)

James Neilson (GDG Chair)

Emeritus Professor of Obstetrics and Gynaecology
Editor, Cochrane Pregnancy and Childbirth Group
University of Liverpool
Liverpool, United Kingdom

Lisa Noguchi

Director, Maternal Newborn Health
Jhpiego
Baltimore, Maryland, United States of America (USA)

Nafissa Osman

Associate Professor of Obstetrics/Gynaecology
Faculty of Medicine
Universidade Eduardo Mondlane
Maputo, Mozambique

Erika Ota

Professor
St Luke's International University
Tokyo, Japan

Robert Pattinson

Director
South Africa Medical Research Center/
University of Pretoria
Maternal and Infant Health Care Strategies
Research Unit
Pretoria, South Africa

Kathleen Rasmussen

Professor of Maternal and Child Nutrition
Cornell University
Ithaca, New York, USA

Rusidah Selamat

Deputy Director (Nutrition Policy and Planning)
Nutrition Division
Ministry of Health
Putrajaya, Malaysia

Harshpal Singh Sachdev

Senior Consultant in Pediatrics and Clinical Epidemiology
Sitaram Bhartia Institute of Science and Research
New Delhi, India

Charlotte Warren

Senior Associate, Maternal and Newborn Health
Population Council
Washington DC, USA

Guideline Methodology Team**Theresa Lawrie**

Director
The Evidence-Based Medicine Consultancy
Bath, United Kingdom

Therese Dowswell

Systematic Reviewer
The Evidence-Based Medicine Consultancy
Bath, United Kingdom

Observers**Nita Dalmiya**

United Nations Children's Fund
New York City, New York, USA

Lindy Fenlason

United States Agency for International
Development (USAID)
Washington DC, USA

Mary Ellen Stanton

USAID
Washington DC, USA

Petra Ten Hoop-Bender

United Nations Population Fund
Geneva, Switzerland

Allison Tumilowicz

Bill & Melinda Gates Foundation
Seattle, Washington, USA

Florence West

International Confederation of Midwives
Hague, Netherlands

External Review Group

Lorena Binfa

Professor
Department of Women's and Newborn Health
Promotion
Faculty of Medicine
University of Chile
Santiago, Chile

Tamar Kabakian

Associate Professor
Health Promotion and Community Health
Faculty of Health Sciences
American University of Beirut
Beirut, Lebanon

Bhavya Reddy

Senior Research Associate
Ramalingaswami Centre on Equity and Social
Determinants of Health
Public Health Foundation of India
Bangalore, India

Petr Velebil

Obstetrician
Perinatal Centre of the Institute for the
Care of Mother and Child
Prague, Czech Republic

Annex 2. Summary of declarations of interest from the Guideline Development Group members, observers and External Review Group members and how they were managed

Name	Gender	Expertise	Disclosure of interest	Conflict of interest and management
Dr Niveen Abu-Rmeileh	Female	Community and public health, statistical epidemiology	None declared	None declared
Professor Lorena Binfa	Female	Midwifery, Director of the WHO collaborating centre for Developing Midwifery	None declared	None declared
Dr Nita Dalmiya	Female	United Nations Children's Fund New York City, New York, United States of America (USA)	None declared	None declared
Dr Lindy Fenlason	Female	United States Agency for International Development	None declared	Not applicable
Dr Atf Ghérissi	Female	Systematic reviews, qualitative evidence, maternal and perinatal health, community health	None declared	Not applicable
Mrs Gill Gyte	Female	Consumer representative, pregnancy and childbirth	None declared	Not applicable
Professor Tamar Kabakian	Female	Health Promotion and Community Health	None declared	None declared
Professor Jim Neilson	Male	General obstetrics, perinatology, gynaecology, systematic reviews, evidence synthesis and guideline development using GRADE	None declared	Not applicable
Dr Lisa Noguchi	Female	Midwifery, delivery of care, implementation science	Employer anticipated research funding from Bill & Melinda Gates Foundation related to studying introduction of innovations and improving quality of care in ANC and PNC.	The conflict was not considered serious enough to affect Guideline Development Group (GDG) membership or participation in the GDG meeting.

Name	Gender	Expertise	Disclosure of interest	Conflict of interest and management
Professor Nafissa Osman	Female	Obstetrics and gynaecology, implementation research	None declared	Not applicable
Professor Erika Ota	Female	Nutrition, evidence synthesis, guideline development	None declared	Not applicable
Professor Bob Pattinson	Male	Obstetrics and gynaecology, delivery of care, evidence synthesis	None declared	Not applicable
Professor Kathleen Rasmussen	Female	Professor of maternal and child nutrition	None declared	Not applicable
Ms Bhavya Reddy	Female	Maternal and child health, gender and health equity, community health	None declared	None declared
Professor Harshpal Singh Sachdev	Male	Paediatrics, nutrition, systematic reviews	Contributed data from India to subsequent meta-analyses and contributed to a published opinion paper on the subject of zincs in pregnancy. Was involved in the epidemiological design and analysis of this paper; however, did not receive funding for this work	The conflict was not considered serious enough to affect GDG membership or participation in the GDG meeting.
Ms Rusidah Selamat	Female	Maternal and infant nutrition, community-based programmes, implementation research	None declared	Not applicable
Dr Mary Ellen Stanton	Female	United States Agency for International Development	None declared	Not applicable
Dr Petra Ten Hoopen-Bender	Female	United Nations Population Fund	None declared	Not applicable
Dr Alison Tumilowicz	Female	Bill & Melinda Gates Foundation	None declared	Not applicable
Dr Petr Velebil	Male	Obstetrics and gynaecology	None declared	None declared
Dr Charlotte Warren	Female	Maternal and perinatal health, systematic reviews, implementation research	None declared	Not applicable
Dr Florence West	Female	International Confederation of Midwives	None declared	Not applicable

Annex 3. Zinc supplementation: GRADE tables

Question: Should zinc supplementation versus no zinc supplementation (with or without placebo) be used for pregnant women?

Settings: Bangladesh, Chile, China, Denmark, Indonesia, Islamic Republic of Iran, United Kingdom of Great Britain and Northern Ireland, United Republic of Tanzania, United States of America.

Bibliography: Carducci B, Keats EC, Bhutta ZA. Zinc supplementation for improving pregnancy and infant outcome. Cochrane Database Syst Rev. 2021; 3:CD000230. doi:10.1002/14651858.CD000230.pub6.

Maternal outcomes

Number of studies	Design	Certainty assessment					Number of women		Effect		Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Zinc supplementation	No zinc (with or without placebo)	Relative (95% CI)	Absolute		
Caesarean section												
7	randomized trials	serious ^a	serious ^b	no serious indirectness	serious ^c	none	150/1132 (13.3%)	143/1079 (13.3%)	RR 0.87 (0.58 to 1.29)	17 fewer per 1000 (from 56 fewer to 38 more)	⊗○○○ VERY LOW	CRITICAL
								31.8%		41 fewer per 1000 (from 134 fewer to 92 more)		
Instrumental vaginal birth												
1	randomized trial	serious ^d	no serious inconsistency	no serious indirectness	serious ^c	none	58/585 (9.9%)	55/621 (8.9%)	RR 1.12 (0.79 to 1.59)	11 more per 1000 (from 19 fewer to 52 more)	⊗⊗○○ LOW	CRITICAL
								8.9%		11 more per 1000 (from 19 fewer to 53 more)		

CI = confidence interval, RR = risk ratio

Number of studies	Design	Certainty assessment					Number of women			Effect		Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Zinc supplementation	No zinc (with or without placebo)	Relative (95% CI)	Absolute			
Pre-eclampsia													
6	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	serious ^c	none	41/1265 (3.2%)	45/1303 (3.5%)	RR 0.93 (0.62 to 1.42)	2 fewer per 1000 (from 13 fewer to 15 more)	RR 0.93 (0.62 to 1.42)	⊗⊗○○ LOW	CRITICAL
								4.4%		3 fewer per 1000 (from 17 fewer to 18 more)			
Any maternal infection													
4	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	serious ^c	none	86/930 (9.2%)	96/961 (10%)	RR 0.94 (0.72 to 1.23)	6 fewer per 1000 (from 28 fewer to 23 more)	RR 0.94 (0.72 to 1.23)	⊗⊗○○ LOW	CRITICAL
								5.4%		3 fewer per 1000 (from 15 fewer to 12 more)			
								18.2%		47 fewer per 1000 (from 116 fewer to 91 more)			

^a Most of the pooled effect was provided by studies with moderate risk of bias and design limitations.

^b Inconsistency between studies contributing data (I² = 72%) (size and direction of effect inconsistent).

^c Confidence interval is imprecise.

^d Evidence from a single study with moderate risk of bias and design limitations.

Fetal and neonatal outcomes

Number of studies		Design	Certainty assessment				Number of women			Effect		Certainty	Importance
			Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Zinc supplementation	No zinc (with or without placebo)	Relative (95% CI)	Absolute		
Small for gestational age or intrauterine growth restriction													
9	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	no serious imprecision	none	609/2695 (22.6%)	590/2635 (22.4%)	RR 1.02 (0.92 to 1.12)	4 more per 1000 (from 18 fewer to 27 more)	⊗⊗⊗○ MODERATE	CRITICAL	
								14%		3 more per 1000 (from 11 fewer to 17 more)			
Low birthweight (< 2500 g)													
17	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	no serious imprecision	none	499/3755 (13.3%)	483/3644 (13.3%)	RR 0.94 (0.79 to 1.13)	8 fewer per 1000 (from 28 fewer to 17 more)	⊗⊗⊗○ MODERATE	CRITICAL	
								8.3%		5 fewer per 1000 (from 17 fewer to 11 more)			
Preterm birth (< 37 weeks)													
20	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	no serious imprecision	none	557/4756 (11.7%)	599/4698 (12.8%)	RR 0.9 (0.78 to 1.04)	13 fewer per 1000 (from 28 fewer to 5 more)	⊗⊗⊗○ MODERATE	CRITICAL	
								8.9%		9 fewer per 1000 (from 20 fewer to 4 more)			

Number of studies	Design	Certainty assessment					Number of women		Effect		Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Zinc supplementation	No zinc (with or without placebo)	Relative (95% CI)	Absolute		
Perinatal death												
2	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	serious ^b	none	81/1272 (6.4%)	70/1217 (5.8%)	RR 1.1 (0.81 to 1.51)	6 more per 1000 (from 11 fewer to 29 more)	⊗⊗○○ LOW	CRITICAL
								5.7%				
Neonatal death												
3	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	serious ^b	none	36/1013 (3.6%)	23/952 (2.4%)	RR 2.44 (0.4 to 14.83)	35 more per 1000 (from 14 fewer to 334 more)	⊗⊗○○ LOW	CRITICAL
								0%				
Stillbirth												
6	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	serious ^b	none	48/1450 (3.3%)	35/1448 (2.4%)	RR 1.32 (0.85 to 2.04)	8 more per 1000 (from 4 fewer to 25 more)	⊗⊗○○ LOW	CRITICAL
								2.3%				

Number of studies	Design	Certainty assessment					Number of women			Effect		Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Zinc supplementation	No zinc (with or without placebo)	Relative (95% CI)	Absolute			
Congenital malformation													
5	randomized trials	serious ^c	no serious inconsistency	no serious indirectness	serious ^b	none	13/595 (2.2%)	18/511 (3.5%)	RR 0.67 (0.33 to 1.35)	12 fewer per 1000 (from 24 fewer to 12 more)	⊗⊗○○ LOW	CRITICAL	
								3.2%		11 fewer per 1000 (from 21 fewer to 11 more)			
Neonatal sepsis													
2	randomized trials	serious ^a	no serious inconsistency	no serious indirectness	serious ^b	none	1/410 (0.24%)	6/326 (1.8%)	RR 0.17 (0.03 to 1.01)	15 fewer per 1000 (from 0 fewer to 18 fewer)	⊗⊗○○ LOW	CRITICAL	
								2.1%		17 fewer per 1000 (from 0 fewer to 20 fewer)			

^a Evidence from studies with moderate risk of bias and design limitations.

^b Confidence interval is imprecise.

^c More than 50% of the pooled effect from studies with moderate risk of bias and design limitations.

^d Estimate based on low event rate.



For more information, please contact:

Department of Sexual and Reproductive Health and Research

Fax: +41 22 791 4171

Email: reproductivehealth@who.int

Website: https://www.who.int/reproductivehealth/about_us/en/

Department of Maternal, Newborn, Child and Adolescent Health, and Ageing

Tel. +41 22 791 3281

Fax: +41 22 791 4853

Email: mncah@who.int

Website: www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing

Department of Nutrition and Food Safety

Fax: +41 22 791 4156

Email: nutrition@who.int

Website: <https://cms.who.int/teams/nutrition-and-food-safety/overview>

World Health Organization
20 Avenue Appia, 1211 Geneva 27
Switzerland

ISBN 978 92 4 003046 6

