
Hypertension Clinical Pathway



ACCURATE BLOOD PRESSURE MEASUREMENT

MEASURE BLOOD PRESSURE IN ALL ADULTS AND AT ALL VISITS







CARDIOVASCULAR RISK

KNOW YOUR RISK OF CARDIOVASCULAR DISEASE AND HOW TO MODIFY IT

CARDIOVASCULAR RISK **CALCULATOR**

Use the **HEARTS** App to assess your cardiovascular risk



Scan code to access the cardiovascular risk calculator



This App does not replace clinical judgment.



TREATMENT PROTOCOL

START TREATMENT IMMEDIATELY AFTER **CONFIRMING HYPERTENSION**

Blood Pressure ≥140/90 mmHg in all HYPERTENSIVES. Systolic Blood Pressure ≥130 mmHg in HIGH-RISK HYPERTENSIVES (Established cardiovascular disease, Diabetes, Chronic Kidney Disease, Risk score ≥

| Cardiovascular risk | | All Hypertensives | HIGH-RISK Hypertensives | | |
|--|----------|----------------------|---|--|--|
| | | | WITH established cardiovascular disease | WITHOUT established cardiovascular disease | |
| Blood Pressure TARGET <140/90 mmHg | | √ | | | |
| Systolic Blood Pressure TARGET <130 mmHg | | | √ | √ | |
| ASPIRIN mg/daily | | | √ | | |
| High-dose statins: | mg/daily | | √ | | |
| Moderate-dose statins: | mg/daily | | | √ | |



Avoid alcohol consumption

Patient above target after repeat measurement



Do 30 minutes of physical activity daily



Body mass index between 18.5 and 24.9



Avoid foods high in sodium

Patient above target after repeat measurement



1 MONTH

1 MONTH

1 MONTH



Patient above target after repeat measurement

1 MONTH



No smoking

| Patient above targ | et: |
|-------------------------|---------|
| Refer to the next level | of care |

| Patients under control | Minimum 6-MONTH follow-up | Minimum 3-MONTH follow-up | Supply medicines for 3 MONTHS | Vaccination | | |
|-------------------------|------------------------------|------------------------------|-------------------------------|-------------|--------------|-------|
| | | | | Influenza | Pneumococcus | COVID |
| All Hypertensives | | | | | | / |
| HIGH-RISK Hypertensives | | | | \ | / | |





ASSESS TREATMENT ADHERENCE AT EACH VISIT

TAKE ALL MEDICATIONS AT THE SAME TIME EVERY DAY

This protocol is **NOT INDICATED** in **WOMEN** of **CHILDBEARING** AGE