



HEARTS IN THE AMERICAS Compendium of Essential Clinical Tools 2023



HE RTS IN THE AMERICAS

HEARTS in the Americas. Compendium of essential clinical tools 2023

PAHO/NMH/NV/24-0002

© Pan American Health Organization, 2024

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO license (CC BY-NC-SA 3.0 IGO).

Under the terms of this license, this work may be copied, redistributed, and adapted for non-commercial purposes, provided the new work is issued using the same or equivalent Creative Commons license and it is appropriately cited. In any use of this work, there should be no suggestion that the Pan American Health Organization (PAHO) endorses any specific organization, product, or service. Use of the PAHO logo is not permitted.

All reasonable precautions have been taken by PAHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall PAHO be liable for damages arising from its use.

HEARTS in the Americas for Better Care for Noncommunicable Diseases (NCDs) in Primary Care Settings

HEARTS in the Americas is an operational framework that will contribute to increasing equitable access to comprehensive NCD management in primary health care (PHC) under PAHO's new initiative "Better Care for NCDs: Accelerating Actions in Primary Health Care."





- Strengthening the capacity of health authorities to plan and implement this approach.
- Increasing the capacity of health services to deliver comprehensive quality screening, diagnosis, treatment, and follow-up.
- Strengthening data collection and monitoring.

Multidisciplinary approach focused on six strategic areas:



HEARTS in the Americas

A platform for expanding comprehensive NCD services









Introduction

Cardiovascular diseases are the leading cause of mortality and morbidity in the Region of the Americas and hypertension is their main risk factor. Despite the remarkable progress made in the last four decades, the observed trend of declining mortality from these diseases has slowed in most countries, and has even started to increase in a group of them. This scenario puts at risk the goal of reducing premature mortality from noncommunicable diseases (NCDs) by 30% by 2030, a commitment made by all countries in the Region.¹

There is a strong association between the level of hypertension control in the population and mortality from cardiovascular diseases. Countries with higher levels of population control of hypertension have achieved greater reductions in mortality from these diseases. In the Americas, for every 1% increase in the level of control of hypertension in the population, there is an estimated 2.9% reduction in deaths from ischemic heart disease per 100 000 population, equivalent to 25 639 avoidable deaths (2.5 deaths per 100 000 population) and 2.37% reduction in deaths from stroke per 100 000 population, equivalent to 9650 avoidable deaths (1 death per 100 000 population). Indeed, if Latin America and the Caribbean increased the level of population control of hypertension from 28.7% to 50%, about 459 000 deaths could be avoided (331 000 from ischemic heart disease and 128 000 from stroke).²

The slowdown in the cardiovascular disease mortality decline in the Americas, combined with suboptimal levels of hypertension control, suggests that the current healthcare services model is exhausted, and a paradigm shift is imperative. In this context, the HEARTS in the Americas Initiative by the Pan American Health Organization (PAHO) was born as a regional adaptation of the World Health Organization (WHO) Global HEARTS Initiative. HEARTS in the Americas will be the institutionalized model for hypertension control and comprehensive cardiovascular risk management throughout the Region by 2025. Based on primary care, this initiative is led by health ministries and supported technically by PAHO. It is currently being implemented in 33 countries in the Region and aims to be fully integrated into each country's healthcare system.³

HEARTS in the Americas has developed several tools to assist healthcare teams in their daily work and encourage them to seek continuous quality improvement to achieve the desired objectives. These clinical tools have been compiled in this document for easy access.

Purpose

The purpose of *HEARTS in the Americas: Compendium of Essential Clinical Tools* is to present, in a simple and instructive way, all the technical resources that the initiative has developed to facilitate the daily work of healthcare teams. The reader will notice that each tool is summarized on a single page, and it is presented in a modular format. Therefore, the tools can be used together or separately as needed.

HEARTS in the Americas proposes this simple, evidence-based, action-oriented compendium, adapted to a digital era in which scientific information overload makes it challenging to access fundamental concepts and where professionals' heavy work load does not allow them sufficient time for updating their knowledge.

The HEARTS in the Americas team hopes that this Compendium of Essential Clinical Tools aims to guide the daily work of healthcare teams and help improve current practices to provide the highest-quality and comprehensive cardiovascular risk management possible and ultimately reduce the burden of cardiovascular disease.

Authorship and acknowledgments

This Compendium of Essential Clinical Tools was prepared by Andres Rosende, International Consultant for the PAHO HEARTS in the Americas Initiative, under the technical coordination of Pedro Ordunez, Regional Advisor for Chronic Diseases and Mental Health at PAHO and technical coordinator of the HEARTS in the Americas Initiative, and with the collaborative work of the Regional HEARTS in the Americas team: Angelo Gamarra, Cintia Lombardi, Natalia Parra, Libardo Rodríguez, and Yenny Rodríguez.

Furthermore, the authors would like to thank Teresa Aumala, Alison Mundaca, Carolina Neira, Michael Pereira, Gonzalo Rodríguez, Yamilé González Valdés, and Eric Zúñiga, all healthcare and HEARTS colaborators, for their critical review and contributions to the improvement of this compendium.

1. Martinez R, Soliz P, Mujica OJ, Reveiz L, Campbell NRC, Ordunez P. The slowdown in the reduction rate of premature mortality from cardiovascular diseases puts the Americas at risk of achieving SDG 3.4: A population trend analysis of 37 countries from 1990 to 2017. J Clin Hypertens (Greenwich). 2020;22(8):1296–1309. Available from: https://doi.org/10.1111/jch.13922.

2. Martinez R, Soliz P, Campbell NRC, Lackland DT, Whelton PK, Ordunez P. Association between population hypertension control and ischemic heart disease and stroke mortality in 36 countries of the Americas, 1990-2019: an ecological study. Rev Panam Salud Publica. 2022;46:e143. Available from: https://doi.org/10.26633/RPSP.2022.143.

3. Ordunez P, Campbell NRC, Giraldo Arcila GP, Angell SY, Lombardi C, Brettler JW, et al. HEARTS in the Americas: innovations for improving hypertension and cardiovascular disease risk management in primary care. Rev Panam Salud Publica. 2022;46:e96. Available from: https://doi.org/10.26633/RPSP.2022.96.











HEARTS in the Americas: a new paradigm in the prevention of cardiovascular disease

The trend toward a reduction in cardiovascular mortality in the Americas has stagnated and even reversed in many countries. HEARTS in the Americas represents a paradigm shift in the comprehensive approach to cardiovascular disease (CVD) prevention, focusing on hypertension diagnosis and treatment as the main risk factor for CVD.



Martinez R, Soliz P, Campbell NRC, Lackland DT, Whelton PK, Ordunez P. Association between population hypertension control and ischemic heart disease and stroke mortality in 36 countries of the Americas, 1990-2019: an ecological study. Rev Panam Salud Publica. 2022;46:e143. Available from: https://doi.org/10.26633/RPSP.2022.143.

Ordunez P, Campbell NRC, Giraldo Arcila GP, Angell SY, Lombardi C, Brettler JW, et al. HEARTS in the Americas: innovations for improving hypertension and cardiovascular disease risk management in primary care. Rev Panam Salud Publica. 2022;46:e96. Available from: https://doi.org/10.26633/RPSP.2022.96.





Protocol for standardized blood pressure measurement

Lack of standardization in blood pressure measurement is a serious and frequent problem in clinical practice that impacts patient safety and health system performance.

3 STEPS FOR ACCURATE BLOOD PRESSURE MEASUREMENT



Before measurement

- Provide a quiet environment at a comfortable temperature.

2.

- Explain the procedure to be performed.
- Check that the patient:
 - Has NOT smoked or ingested food, coffee, or alcohol in the last 30 MINUTES
 - Has had at least **5 MINUTES** of rest.

USE A CLINICALLY VALIDATED BLOOD PRESSURE MEASURING DEVICE

During measurement

Ensure these 8 conditions are met





✓ Obtain a SECOND MEASUREMENT with an interval of 30 SECONDS

Register both **MEASUREMENTS WITHOUT ROUNDING** and calculate the **AVERAGE** to obtain the **FINAL RESULT**

Cheung AK, Whelton PK, Muntner P, Schutte AE, Moran AE, Williams B, International Consensus on Standardized Clinic Blood Pressure Measurement - A Call to Action. Am J Med. 2023;136(5):438–445.e1. Available from: https://doi.org/10.1016/j.amjmed.2022.12.015.

Access a video on blood pressure measurement by clicking here:





Resources to verify if a blood pressure measuring device is clinically validated

Clinical validation refers to the rigorous testing of blood pressure measuring devices for accuracy in clinical practice. It is essential to obtain reliable measurements that allow making appropriate therapeutic decisions, increasing patient safety. The table below includes lists of blood pressure measuring devices that have passed validation studies. Instructions on how to use these lists are shown in the reference underneath.

Websites containing lists of validated automated blood pressure measuring devices

| STRIDE BP | STRIDE BP is an international scientific nonprofit organization founded by hypertension experts with the mission of improving the accuracy of blood pressure measurement and the diagnosis and management of hypertension. STRIDE BP provides international guidance and practice tools on the methodology and technology for accurate blood pressure evaluation according to the latest scientific evidence. STRIDE BP is cosponsored by the European Society of Hypertension and the International Society of Hypertension. | |
|----------------|---|--|
| Medaval | Medaval is a company that assesses and lists both validated and nonvalidated blood pressure devices available globally. It has the largest database. Assessment of validation status is performed according to international scientific protocols. Certification of validation according to current accuracy assessment protocols is provided by Medaval on application and payment by manufacturers. Validation and comparative-equivalence reports are peer reviewed by members from its reviewer panels. It also provides links to the recommendations of other validated device listings. | |
| US Blood | An independent process was designed to determine which blood pressure | |
| Pressure | devices available in the United States meet the American Medical Association | |
| Validated | (AMA) established criteria to validate clinical accuracy. This review process | |
| Device Listing | results in a formal list of blood pressure devices (the "Validated Device Listing" | |
| (VDL) | or "VDL"). AMA does not receive funding from manufacturers. | |
| British and | All monitors on the "Home" and "Specialist Use" lists have been approved | |
| Irish | by BIHS for accuracy. The list also includes nonvalidated devices. | |
| Hypertension | Those bearing the BIHS logo have been tested in-house. The rest have | |
| Society | been tested by independent agencies, and their published reports have been | |
| (BIHS) | peer-reviewed and approved by BIHS. | |

*PAHO does not endorse these lists or the devices included in them.

3

Picone DS, Padwal R, Campbell NRC, Boutouyrie P, Brady TM, Olsen MH, et al. How to check whether a blood pressure monitor has been properly validated for accuracy. J Clin Hypertens (Greenwich). 2020;22(12):2167–2174. Available from: https://doi.org/10.1111/jch.14065.





HEARTS App: a pragmatic approach to estimating cardiovascular disease risk

Estimation of cardiovascular risk is essential to optimize healthcare system interventions. HEARTS in the Americas proposes a simple and pragmatic approach, based on the WHO hypertension guidelines, to effectively guide clinical practice. The HEARTS App is based on the following algorithm. It includes a calculator derived from the WHO cardiovascular risk tables and contains the clinical pathways defined by each country.



- achieved
- Follow up at least every 3 months if BP is under control

Note: BP: blood pressure; CVD: cardiovascular disease; CKD: chronic kidney disease; SBP: systolic blood pressure; DBP: diastolic blood pressure

Ordunez P, Tajer C, Gaziano T, Rodriguez YA, Rosende A, Jaffe MG. The HEARTS app: a clinical tool for cardiovascular risk and hypertension management in primary health care. Rev Panam Salud Publica. 2022;46:e12. Available from: https://doi.org/10.26633/RPSP.2022.12.





5.

The 2021 WHO guideline for the pharmacological treatment of hypertension in adults

The WHO hypertension guideline is a methodologically rigorous, evidence-based document with broad international consensus focusing on implementation.

8 recommendations in 5 questions



When **TO INITIATE** pharmacological **TREATMENT**?



R1. BP thresholds to initiate medication

- ✓ Those with a diagnosis of hypertension and BP ≥140/90 mmHg
- Those with CVD and SBP 130–139 mmHg

Recommendation: **strong** Evidence: **moderate-high certainty**

 Those without CVD but with HIGH CVD RISK and SBP 130–139 mmHg

Initiate pharmacological treatment within 4 WEEKS of diagnosis

Recommendation: conditional Evidence: moderate-high certainty



What are the control TARGETS?

R6. BP control targets

- ✓ BP <140/90 mmHg in those WITHOUT HIGH CVD RISK</p>
- ✓ SBP <130 mmHg in those WITH CVD</p>

Recommendation: **strong** Evidence: **moderate certainty**

 SBP <130 mmHg in those without CVD but with HIGH CVD RISK

Recommendation: conditional Evidence: moderate certainty



How to establish the FOLLOW-UP intervals?



What MEDICATIONS to use?



R4. First-line medications

- ✓ ACEi/ARB
- Dihydropyridine CCB
- TZ/TZ-like diuretic

Recommendation: **strong** Evidence: **high certainty**



R5. Combined treatment

COMBINE an ACEi or ARB + a CCB or a TZ/TZ-like diuretic

Preferably in a **SINGLE PILL COMBINATION** to improve adherence and persistence

Recommendation: conditional Evidence: moderate certainty



When to assess CVD RISK?

- **R2 & 3. CVD RISK**
- Obtain laboratory tests and assess CVD risk systematically but only if IT DOES NOT DELAY HYPERTENSION TREATMENT

Recommendation: conditional Evidence: low certainty

R7. Follow-up intervals



✓ MONTHLY until reaching BP CONTROL

EVERY 3–6 MONTHS in CONTROLLED BP

Recommendation: conditional Evidence: low certainty



 TREATMENT PROVIDED by NONPHYSICIAN PROFESSIONALS under supervision and following a protocol

Recommendation: **low certainty** Evidence: **conditional**

Note: ACEi: angiotensin converting enzyme inhibitor; ARB: angiotensin receptor blocker; CCB: calcium channel blocker; TZ: thiazide

Adapted from World Health Organization. Guideline for the pharmacological treatment of hypertension in adults: summary. Geneva: WHO; 2022. Available from: https://iris.who.int/handle/10665/356108.

Campbell NRC, Paccot Burnens M, Whelton PK, Angell SY, Jaffe MG, Cohn J, et al. 2021 World Health Organization guideline on pharmacological treatment of hypertension: Policy implications for the region of the Americas. Lancet Reg Health Am. 2022;9:100219. Available from: https://doi.org/10.1016/j.lana.2022.100219.





Key interventions for the management of cardiovascular disease risk

Improving clinical processes is a central element in closing quality gaps in the delivery of health services and positively impacting coverage and control indicators. For this reason, the HEARTS Innovation Group has defined key interventions to achieve this.

| 5 Areas | 8 Drivers | Interventions |
|-----------------------|--------------------------------|--|
| Diagnosis | Accuracy of BP measurement | a. Blood pressure measurement training every 6 months b. Blood pressure measurement protocol c. Use automated and clinically validated BPMDs |
| Blagnoolo | CVD risk assessment | a. Classify patients according their CVD risk b. Use statins and aspirin according to risk level and history of CVD |
| Treatment | Treatment protocol | a. Institutionalized clinical pathwayb. Use fixed dose combinations |
| | Treatment intensification | a. Initiate treatment without delaysb. Intensify treatment until reaching BP control |
| Continuity of care | Follow-up frequency | a. Monthly follow-up until reaching BP control b. 6-monthly follow-up in non high-risk patients with controlled BP c. 3-monthly follow-up in high-risk patients with controlled BP |
| Delivery system | Team-based care | a. Non physician community health workers measure BP b. Nurses follow patients c. Nurses intensify pharmacological treatment following the clinical pathway |
| | Medication refill frequency | a. Medication refill every 3 months |



Note: BPMDs: blood pressure measuring devices

6

Brettler JW, Arcila GPG, Aumala T, Best A, Campbell NR, Cyr S, et al. Drivers and scorecards to improve hypertension control in primary care practice: Recommendations from the HEARTS in the Americas Innovation Group. Lancet Reg Health Am. 2022;9:100223. Available from: https://doi.org/10.1016/j.lana.2022.100223.





7.

The HEARTS in the Americas Clinical Pathway

The clinical pathway represents the main implementation tool of the HEARTS in the Americas Initiative. It summarizes, in a single page, the comprehensive management of cardiovascular risk in primary health care, incorporating the WHO hypertension guideline recommendations and the HEARTS drivers. Hence, the clinical pathway becomes a powerful tool to strengthen management, clinical practice, and community action.



Rosende A, DiPette D, Brettler J, Rodríguez G, Zuniga E, Connell K, et al. HEARTS in the Americas appraisal checklist and clinical pathway for comprehensive hypertension management in primary care. Rev Panam Salud Publica. 2022;46:e125. Available from: https://doi.org/10.26633/RPSP.2022.125.

Rosende A, DiPette DJ, Martinez R, Brettler JW, Rodriguez G, Zuniga E, et al. HEARTS in the Americas clinical pathway. Strengthening the decision support system to improve hypertension and cardiovascular disease risk management in primary care settings. Front Cardiovasc Med. 2023;10:1102482. Available from: https://doi.org/10.3389/ fcvm.2023.1102482.





Monitoring and Evaluation System for quality improvement

The System for Monitoring and Evaluation of HEARTS in the Americas is a tool for quality management based at the primary health care center. Developed under the open-source platform DHIS2, it allows health teams to report the variables linked to their professional practice and obtain periodic reports that enable them to evaluate their performance and results.



Based on the health center

- Designed for and by the primary health care team.
- Real-time analytical and systematized report.
- Control dashboard to assess implementation maturity and performance over time.

Built on the open-source DHIS2 platform

- Interoperable with multiple electronic health record systems. \checkmark
- Free, open for use, and does not require a permanent Internet connection. \checkmark
- Dynamic technology adoption, ensuring robust solutions that operate in \checkmark resource-constrained environments.





Indicators aligned with programmatic priorities

- Aimed at evaluating all aspects of HEARTS implementation.
- Aggregate data to protect the privacy of the medical record.
- Allows for the identification of implementation gaps.



Instrumental for the continuous quality improvement process

- Access to information under a hierarchical structure to facilitate management at different levels.
- Allows comparison between different health centers to promote learning based on best practices.
- Generates evidence to guide decision-making, identify common problems, and support strategies for target-oriented intervention design.



Prado P, Gamarra A, Rodriguez L, Brettler J, Farrell M, Girola ME, et al. Monitoring and evaluation platform for HEARTS in the Americas: improving population-based hypertension control programs in primary health care. Rev Panam Salud Publica. 2022;46:e161. Available from: https://doi.org/10.26633/RPSP.2022.161.





9.

HEARTS training and education resources

Continuous training and education of the healthcare team is essential to provide quality care. HEARTS in the Americas has developed a set of free and open access courses with the collaboration of scientific organizations and international experts. Each course provides a PAHO certificate with academic hours and a unique badge.



Hypertension Control Drivers at Primary Health Care Centers



Implementation of the HEARTS Technical Package in Primary Care Health Teams



Accurate Automated Blood Pressure Measurement



Supporting Primary Health Care Teams to Use Blood Pressure Medications Effectively







Hypertension and Cardiovascular Risk Management in Primary Health Care

Home Blood Pressure Monitoring

Access HEARTS virtual courses by clicking here:





Since 2016, the Pan American Health Organization (PAHO) has been promoting the implementation of the HEARTS Initiative in the Americas as a regional adaptation of the World Health Organization's Global HEARTS Initiative. During this time, 33 countries in the Region of the Americas have committed to implementing HEARTS, and PAHO has developed many technical resources to support them. Most of these resources are clinical tools for primary healthcare teams and focus on quality improvement.

This compendium aims to group all these clinical tools in a single document under a simple format that facilitates their practical implementation in daily clinical practice. The reader will notice that each tool is summarized on a single page and presented in a modular format. Therefore, each tool can be used together or separately as needed. In addition, at the bottom of each tool, the reader will find the references and hyperlinks to access full texts in case deeper knowledge is required.



